

Table S2. Pairwise geographic distances (in Km, below diagonal) and genetic differentiation (Slatkin's F_{ST} , above diagonal) among *Sorex minutus* phylogroups and other geographic groups

	Italian	South Italian	Balkan	Iberian	Belle Île	Britain	Northern (Continental)	Western (Continental)	Orkney Islands	Irish
Italian	-	1.6558	3.0387	3.6919	1.594	2.8852	2.3798	1.4534	2.8673	4.009
South Italian	773.14	-	2.5113	3.3869	1.3569	2.8562	2.3204	1.1713	2.7079	4.682
Balkan	547.27	628.96	-	1.9234	2.5617	3.1494	2.8093	2.7975	6.2456	7.0533
Iberian	1349.26	1768.56	1880.98	-	4.265	3.185	2.6191	3.8498	7.3804	10.8797
Belle Ile	1162.82	1815.47	1701.44	640.58	-	2.7595	2.179	0.3345	1.2148	2.4003
Britain	1347.34	2108.66	1795.12	1286.79	647.11	-	0.1449	2.6083	3.9225	4.7265
Northern (Continental)	1022.36	1488.84	863.37	2218.78	1788.28	1554.91	-	2.0767	2.791	3.2035
Western (Continental)	903.9	1444.78	1448.34	476.18	434.81	1006.85	1742.75	-	0.5436	1.1193
Orkney Islands	3476.34	3693.42	3127.67	4679.14	4175.54	3739.02	2477.61	4206.5	-	1.4635
Irish	1652.35	2396.73	2124.06	1311.98	726.86	346.14	1897.29	1151.96	4016.56	-