

Introduction

Reporting by radiographers is now widely adopted in England with a current expansion into cross sectional imaging reporting to support service delivery driven by department of health skills mix initiatives¹.

Factors influencing role development in CT head reporting include the national stroke imaging guidelines², NICE head injury guidelines³, and the national radiologist shortage⁴. Supported by Royal College of Radiologist and the Society and College of Radiographers team working guidance⁵ and case studies of CT head reporting implementation in NHS trusts^{6,7}.



Radiographers are now reporting CT head examinations in at least 17 sites in the UK⁸ and NHS service improvement guidance suggests this could increase in the future⁹.

Aims

To analyse the objective structured examination (OSE) results of the first four cohorts of radiographers (n=23) who successfully completed the postgraduate programme (accredited by the College of Radiographers) in reporting of CT head examinations.




Method

Examinations only included in the OSE where there was agreement between the reports of 3 consultant radiologists.

25 CT head examinations included in OSE – Typical cases are listed below.

Radiographic appearances / pathologies included

Acute Subdural Hematoma
Acute on Chronic Subdural Hematoma
Chronic Subdural Hematoma
Acute Extradural Hematoma
Subarachnoid Haemorrhage
Acute Intracerebral Hematoma
Acute Intraventricular Haemorrhage
Contusion
Mass (solitary, multiple, cavitating and/or eroding – various sites)
Glioma
Meningioma
Metastasis
Aneurysm
Acute Infarction
Chronic Infarction



Associated Findings including:

Mass effect, midline shift, herniation, fracture, sulci effacement

Normal Variants /incidental findings including:

Ischaemic vessel disease, benign calcification, cyst, craniotomy

Inclusion / Marking criteria for OSE

Prevalence of abnormal cases = 50%;
Images included of patients referred from A/E, OP, IP and GP sources;
Wide range of clinical indications included;
Expected answers agreed with External Examiner (Consultant Radiologist);
Candidates indicated if appearances were NORMAL or ABNORMAL and;
provided key details of abnormal appearances and pathology demonstrated;
Sensitivity and specificity calculated using NORMAL / ABNORMAL decision;
Maximum of 5 marks (fractionated) allocated per abnormal case;
Agreement percentage calculated using expected agreed answer.

Results

Individual Cohorts

%	2007-08	2008-09	2011-12	2012-13
Sensitivity	100	97.7	100	100
Specificity	96.6	93.7	100	92.2
Agreement	87.5	90.1	91.7	93.3


Most frequent interpretative errors (In descending order)

Perifocal Oedema when normal for age
Raised Intracranial Pressure when normal for age
Lacunar Infarction when normal for age
Periventricular Small Vessel Disease when normal for age
Traumatic Hematoma as Haemorrhagic extension
Subcortical Ischemia when normal for age
Ventriculomegaly for volume effect
Subdural Hygroma as a Subdural Hematoma
Basal Ganglia Ischemia for Perivascular Space
Cerebral Stroke as Cerebral Tumour

Mean OSE Scores 2007-13 Combined

Students n = 23	Mean (%)	95% CI	SD
Reports n = 575			
Sensitivity	99.3	97.4-99.8	5.65
Specificity	95.6	93.1-97.7	2.89
Agreement	90.7	88.1-90.8	7.14

Mean OSE Scores 2007-13 Combined (%)



Discussion

At the end of this accredited postgraduate programme of study, the radiographers have demonstrated high levels of sensitivity, specificity and agreement over 90% on all measures.

Previous studies^{10,11}, investigating variation between experienced radiologists in the interpretation of CT head examinations, demonstrated agreement rates of 86.6% (13.4 - 20.2% disagreement for major significant abnormalities).

Further work is also needed to confirm the clinical application of these initial encouraging findings, which suggest that more radiographers may be able to contribute to this aspect of the reporting service.

References

- Department of Health (2003) *Radiography Skills mix: a report on the four-tier service delivery model*. London: DOH Stationary Office.
- Department of Health (2008) *Implementing the national stroke service strategy – an imaging guide*. London: DOH Stationary Office.
- National Institute for Health and Care Excellence (2014) *Head injury: Triage, assessment, investigation and early management of head injury in children, young people and adults*. CG176. London: National Institute for Health and Care Excellence.
- The Royal College of Radiologists (2012) *Clinical Radiology UK Workforce Report 2011*. London, the Royal College of Radiologists.
- The Royal College of Radiologists and the Society and College of Radiographers (2012) *Team working in clinical imaging*. London: The Royal College of Radiologists and the Society and College of Radiographers.
- Beale, A. Brown, V. (2011) How to implement best practice in stroke and TIA imaging. *Synergy News*; November:15-17.
- NHS Improvement. (2011) *Diagnostics and stroke-a guide to achieving changes in imaging services to support new stroke pathways*. Available Online from: <https://www.improvement.nhs.uk>; [accessed 12.03.14]
- Society and College of Radiographers (2012) *Scope of radiographic practice survey*. London; SCOR.
- Department of Health (2010) Spending review: local government implications.
- McCarron, M.O., Sands, C., McCarron, p. (2010) *Neuroimaging reports in a general hospital: Results from a quality-improvement program*. *Clinical Neurology and Neurosurgery*, 112(1), pp.54-58.
- Briggs, G.M. Flynn, P.A. Worthington, M. Rennie, L. McKinstry, C.S. (2008) 'The role of specialist neuroradiology second opinion reporting: is there added value?' *Clinical Radiology*, 63, pp791-795.