

Evidence for a gender effect in the impact of intimate partner violence victimization on health-related quality of life

Costa D^{1,2},

Soares J^{3,4},

Lindert J⁵,

Hatzidimitriadou E⁶,

Sundin O⁷,

Toth O⁸,

Ioannidi-Kapolo E⁹,

Degomme O¹⁰,

Cervilla J¹¹,

Barros H^{1,2}.

*dmcosta@med.up.pt

www.ispup.up.pt

¹Dept Clinical Epidemiology, Predictive Medicine and Public Health University of Porto Medical School, Porto, Portugal

²Institute of Public Health of the University of Porto, Porto, Portugal

³ Department of Public Health Sciences, Karolinska Institute, Stockholm, Sweden

⁴ Institution for Health Sciences, Department of Public Health Sciences, Mid Sweden University, Sundsvall, Sweden

⁵ Department of Public Health, Protestant University of Applied Sciences Ludwigsburg, Ludwigsburg, Germany and Department for medical Psychology and Medical Sociology, University of Leipzig, Leipzig, Germany

⁶ Faculty of Health and Social Care Sciences, Kingston University and St George's, University of London, London, United Kingdom

⁷ Division of Psychology, Department of Social Sciences, Mid Sweden University, Östersund, Sweden

⁸ Institute of Sociology, Hungarian Academy of Sciences, Budapest, Hungary

⁹ Department of Sociology, National School of Public Health Athens, Athens, Greece

¹⁰ International Centre for Reproductive Health (ICRH), Ghent University, Ghent, Belgium

¹¹ Departmental Section of Psychiatry and Psychological Medicine, University of Granada, Granada, Spain

Funding:

This work was supported by the Executive Agency for Health and Consumers – European Commission [contract: 20081310] and by the Portuguese Foundation for Science and Technology: SFRH/BD/66388/2009



INTRODUCTION

The effect of intimate partner violence (IPV) on Health-related Quality of Life has been extensively studied in female victims. However, though males are also frequent victims of IPV no such information is available.

AIM

To test in a sample of the European general population the presence of gender differences on how IPV impacts health-related quality of life.

METHODS


Random samples of non-institutionalized adult (18-64 years) men (n=1646) and women (n=2233), residing in 8 European cities (Granada, Gent, Porto, Östersund, Stuttgart, London, Athens and Budapest) were evaluated. Violent acts assessed by the Revised Conflict Tactics Scales were coded to provide past year prevalence for each type of violence (psychological, physical, sexual coercion and injury). When all items were answered with 0 (“This has never happened”), participants were coded as non-cases, and as a case otherwise. A variable was coded to express the severity level (as described by the original scale author in minor and severe acts) for all types of violent sustained in the past year, and used as independent variable: no violence, minor acts, severe acts.

The eight original dimensions of the SF-36 were coded and t-scored to range from 100 to zero (mean=50, standard deviation=10).

The mental (MCS-36) and physical component summary (PCS-36) scores were derived from the eight scales of the SF-36 using principal components analysis with varimax rotation. Higher scores represent better health-related quality-of-life.

General linear models were computed to estimate marginal means of summary components extracted from the SF-36 by violence severity and to test sex interactions. Models were adjusted for country, age and education. Differences in the impact of IPV on the SF-36 summary component scores (Mental and Physical) were expressed as unstandardized betas (Standard Error) according to gender.

European cities represented in the study



Sample size:


- Granada: 138
- Gent: 245
- Porto: 632
- Östersund: 594
- Stuttgart: 546
- London: 571
- Athens: 548
- Budapest: 604

RESULTS

General Linear Model for summary components of the SF-36 by violence severity in the past year

		Mental Health		p-value for sex interaction	Physical Health		p-value for sex interaction
		Women Beta (SE)	Men Beta (SE)		Women Beta (SE)	Men Beta (SE)	
Model 0	No violence	Ref	Ref	<0.001	Ref	Ref	<0.001
	Minor	-1.525 (0.49)	-1.574 (0.54)		1.146 (0.50)	1.263 (0.53)	
	Severe	-5.672 (0.62)	-2.994 (0.62)		-0.530 (0.62) ^{ns}	1.186 (0.62) ^{ns}	
Model 1	No violence	Ref	Ref	<0.001	Ref	Ref	<0.001
	Minor	-1.485 (0.50)	-1.563 (0.54)		0.940 (0.50) ^{ns}	1.199 (0.53)	
	Severe	-5.915 (0.63)	-3.051 (0.64)		-0.567 (0.63) ^{ns}	1.193 (0.64) ^{ns}	
Model 2	No violence	Ref	Ref	<0.001	Ref	Ref	<0.001
	Minor	-1.210 (0.51)	-1.622 (0.54)		-0.100 (0.48) ^{ns}	0.735 (0.50) ^{ns}	
	Severe	-5.511 (0.64)	-2.924 (0.65)		-1.230 (0.61)	0.431 (0.60) ^{ns}	
Model 3	No violence	Ref	Ref	<0.001	Ref	Ref	<0.001
	Minor	-1.049 (0.51)	-1.293 (0.55)		0.076 (0.50) ^{ns}	0.787 (0.52) ^{ns}	
	Severe	-4.182 (0.66)	-1.772 (0.68)		-0.917 (0.65) ^{ns}	0.732 (0.64) ^{ns}	

ns= non-significant; Model 1: adjusted for country; Model 2: adjusted for country, age and education; Model 3: Adjusted for country, age, education and exposure to child abuse.



CONCLUSION

Intimate partner violence influences health-related quality of life in men and women, the impact being more pronounced in the Mental Health of women.