

Research Space

Journal article

Isometric exercise versus high intensity interval training for the management of blood pressure: A systematic review and meta-analysis

Edwards, J., De Caux, A., Donaldson, J., Wiles, J. and O'Driscoll, J.

This is the accepted version of:

Edwards J, De Caux A, Donaldson J, Wiles J, O'Driscoll J. Isometric exercise versus high-intensity interval training for the management of blood pressure: a systematic review and meta-analysis. *Br J Sports Med*. 2021 Dec 15:bjsports-2021-104642. doi: 10.1136/bjsports-2021-104642. Epub ahead of print. PMID: 34911677.

Isometric exercise versus high intensity interval training for the management of blood pressure: A systematic review and meta-analysis.

Edwards, J.J¹., De Caux, A¹., Donaldson, J.A^{1,2}., Wiles, J.D¹., & O'Driscoll, J.M¹.

¹ School of Psychology and Life Sciences, Canterbury Christ Church University, Kent, CT1 1QU

²Department of Sport Science, Nottingham Trent University, Nottinghamshire, NG1 4FQ

Correspondence to Dr Jamie O'Driscoll, School of Psychology and Life Sciences, Canterbury Christ Church University, North Holmes Road, Canterbury, Kent, CT1 1 QU. Email: jamie.odriscoll@canterbury.ac.uk; Telephone: 01227922711.

Running Title: IET vs HIIT in the management of blood pressure

Key Words: Isometric exercise training, high intensity interval training, blood pressure, hypertension

Abstract

Objective: We aimed to compare the efficacy of isometric exercise training (IET) versus high-intensity interval training (HIIT) in the management of resting blood pressure (BP).

Design: Systematic review and meta-analysis.

Data sources: PubMed (MEDLINE), the Cochrane library and SPORTDiscus were systematically searched.

Eligibility criteria: RCT's published between January 1st, 2000 and September 1st, 2020. Research trials reporting the effects of IET or HIIT on resting BP following a short-term intervention (2-12 weeks).

Results: IET produced significantly greater reductions in resting BP compared to HIIT with systolic, diastolic and mean BP effect sizes of 8.50mmHg vs 2.86mmHg (Q=17.10, p<0.001), 4.07mmHg vs 2.48mmHg (Q=4.71, p=0.03) and 6.46mmHg vs 3.15mmHg (Q=4.21, p=0.04) respectively. However, HIIT reduced resting heart rate significantly more than IET (3.17bpm vs 1.34bpm, Q=7.63, p=0.006).

Conclusion: While both modes are efficacious, IET appears to be the superior mode of exercise in the management of resting BP. However, HIIT may achieve wider physiological benefits, with greater reductions in resting heart rate.

Introduction

As the leading modifiable risk factor for mortality, arterial hypertension remains a global problem with important implications for cardiovascular health [1]. Although largely efficacious, anti-hypertensive medication has a number of substantial limitations including adverse side effects [2], considerable economic strain [3] and poor adherence rates, typically reported at less than 50% 1 year following prescription [4]. Additionally, current exercise guidance is unlikely to benefit long-term cardiovascular risk [5] due to poor global adherence [6]. Therefore, establishing novel lifestyle approaches to the management of hypertension is pivotal.

In recent years, isometric exercise training (IET) and high intensity interval training (HIIT) have emerged as convenient and highly time-efficient modes of exercise, which can produce clinically significant reductions in resting clinic and ambulatory blood pressure (BP) following short-term interventions [7–10]. This is crucial as behavioural psychology research has identified perceived lack of time and motivation as the most common barriers to the adoption of physical activity [11]. Importantly, both IET and HIIT interventions have previously been shown to produce a magnitude of BP reduction similar to, or greater than the current recommended international physical activity guidelines [12,13]. Thus, given the efficacy and potential for greater adoption and adherence, these novel exercise modes may have significant clinical utility in the control of arterial BP.

Despite this, no research to date has attempted to compare the anti-hypertensive efficacy of these two distinct exercise modes. Furthermore, previous investigations have only examined the efficacy of HIIT against moderate intensity continuous training or as a sub-group to separate primary analyses, resulting in the omission of a plethora of comprehensive trials and thus an insufficient understanding of the overall effects of HIIT on BP. Additionally, several new IET studies have been published since the latest meta-analysis [14], many of which utilise a lower-body methodology, establishing a need for a lower vs upper body IET analysis. Therefore, we aimed to perform an independent meta-analysis on randomised-controlled trials investigating the effects of IET on BP, with subgroup analysis of the implemented IET method. Separately, we also aimed to perform an independent meta-analysis measuring the effects of HIIT on BP. Finally, following both independent analyses, we performed a meta-analytic comparison to measure the efficacy of IET versus HIIT in the management of BP.

Methods

Search strategy

This systematic review and meta-analysis was performed in accordance with the PRISMA guidelines. To identify potential studies, a systematic search was performed using PubMed (MEDLINE), the Cochrane library and SPORTDiscus for research trials reporting on the effects of IET or HIIT on BP. The search strategy was developed by a research librarian experienced in scientific database searching and included MeSH terms, key words, and word variants. The IET key Boolean search terms included: “Isometric training” OR “Isometric Exercise” OR “Static contraction” OR “Exercise training” AND “Blood pressure” OR Hypertension. The HIIT key Boolean search terms included: “High intensity interval training” OR HIIT OR “anaerobic training” OR “Sprint training” OR “Exercise training” AND “Blood pressure” OR Hypertension. Journal articles written in English and published between the 1st January 2000 and 1st September 2020 were considered. Where possible, corresponding study authors were contacted to ascertain whether non-published data was available or in the pre-print stage, and studies found through the systematic protocol were screened for exterior citations and their respective reference lists searched for eligible studies.

Study eligibility

Screening was carried out following the exclusion of duplicate articles by two authors (JE and AD) who resolved any confliction via consensus. Studies were initially screened by title and then by abstract for relevance. Studies retained for the next step of evaluation were assessed by full-text and considered if they reported pre-post intervention BP changes following either an IET intervention or a HIIT intervention with a control group and sufficient randomisation. Only studies of intervention groups performing IET individually or HIIT individually with no other concurrent exercise were included; however, research designs utilising moderate intensity continuous training as a control arm were included. HIIT was defined in accordance with the EXPERT tool [15], in which >89% heart rate maximum or >80% VO₂ peak was considered a sufficient intensity. Randomised controlled trials, including cross-over design studies with an intervention duration of 2 to 12 weeks were eligible. Participants were required to be ≥18 years of age with no limitations on health or baseline BP status.

Study quality

Study quality and risk of bias was evaluated using the TESTEX scale [16] which is a 15 point (12 item) system designed for the specific application to exercise science research. Two reviewers (JE and AD) independently scored all eligible articles. When disputes were detected in quality analyses, the reviewers met to discuss any conflicts and a third reviewer was consulted (JOD).

Statistical analysis

Derivation of outcome statistics: Data were extracted from the studies independently by the first author. The key variables were 4 continuous measures of cardiovascular health before and after the study intervention; systolic BP (sBP), diastolic BP (dBP), mean BP (mBP) and heart rate with SDs. Where available we also extracted the exercise intervention duration, percentage of medicated participants, and percentage of hypertensive participants for use in the meta-regression analyses. If these data were not available, efforts were made to contact the authors of the paper to obtain them.

Pooling of results: Data were analysed with the meta [17] packages in the statistical software R 4.0.3 (R Core Team, 2020). Effect sizes were calculated based on the change score (MD) between baseline and follow-up measures to demonstrate the clinical impact of each exercise type. Additionally, it has been suggested [18] that change score provides a greater degree of accuracy when the baseline and follow-up scores are highly correlated and SDs are similar, as was the case in our dataset. The MDs with 95% CI's were computed individually for the four outcome measures. The effect sizes were weighted according to the intra-study variability (calculated from reported means and SDs), with larger, more precise studies having a greater weighting on the overall effect size. Separate random-effects meta-analysis were run for each variable using CMA's meta-regression function to establish if any moderator variables influenced the change in BP following IET or HIIT and explain any inter-study variance in outcomes. The moderators assessed independently were: hypertension diagnosis, medication status, TESTEX score, and intervention duration. This yielded the summary measures, significance levels (P), and the between-study heterogeneity (Cochran's Q and I^2) and variability (τ^2).

Exploration of publication bias: Publication bias was assessed via funnel plot inspection, with the expectation that studies would be evenly distributed between both sides of the average effect size, with high-precision studies close to the mean. The individual funnel plots are displayed in the supplementary material (Figure S1). Quantitative assessments of Egger's regression are reported for the four effect sizes alongside the funnel plots.

Exploration of heterogeneity and meta-regression analyses: Heterogeneity of the 3 effect sizes were assessed by Cochran's Q and I^2 values calculated in the random-effects model. Statistical significance for Q was $p < 0.01$ [19] and I^2 of 25, 50 and 75% were interpreted as small, medium and large degrees of heterogeneity. In the event of significant levels of heterogeneity in any of the meta-analyses, exploratory meta-regression analyses were conducted to determine how participant and exercise characteristics accounted for heterogeneous effect sizes. The residual plots were inspected for suitability before reporting the R^2 value of the fitted models. In the event of non-random residual plots, further variables were added to the model to capture possible significant interactions. All covariates were analysed separately and then together to assess the individual contribution of each variable. A method of moment's estimator for the between-study covariance matrix in the random effect models meta-regression was used. A sub-group analysis was conducted based on the MD between study exercise-mode, and 2 groups were formed, which were IET and HIIT. On the IET data we performed additional sub-group analyses for the 4 outcome measures on upper vs. lower body IET training.

Sensitivity analysis: We conducted a sensitivity analysis by removing outlying studies using the leave-one-out influence analysis method in the dmetar package [20]. The study by May et al was found to contribute the most to the overall heterogeneity for all 4 outcome measures but did not influence the overall effect size to a large degree and was left in the final analysis.

Results

As presented in Figure 1, our initial search identified 5220 manuscripts (2793 IET and 2427 HIIT). Following all exclusions, 38 studies were analysed (18 IET and 20 HIIT), including 1583 (672 IET and 911 HIIT) participants, of which 612 (268 IET and 344 HIIT) were controls. Study characteristics for the included IET and HIIT articles can be found in Table 1 and Table 2 respectively. Detailed TESTEX scoring for each category and the full reference list of all analysed studies are presented in the supplementary file (Table S1 and Table S2).

Isometric exercise training

IET significantly reduced sBP and dBP by a mean difference of 8.50mmHg (95% CI= 6.49 to 10.52, $p<0.001$) and 4.07mmHg (95% CI= 3.04 to 5.10, $p<0.001$) respectively, with no significant changes observed in the control group. mBP also significantly decreased following IET by 6.46mmHg (95% CI= 4.18 to 8.74, $p<0.001$) with no significant change in the control group. There were significant differences between groups for all BP variables (all $p<0.001$). IET also had a significant effect on RHR by a mean difference of 1.34bpm (95% CI= 0.24 to 2.44, $p=0.0169$) with no changes in the control group and a significant between-groups effect ($p=0.006$). The subgroup analyses for the effect of upper vs. lower body isometric training were not significant for any of the four outcome measures (sBP, dBP, mBP and RHR).

There was significant heterogeneity between studies in the IET sBP ($I^2=78\%$), dBP ($I^2=52\%$) and mBP arms ($I^2=84\%$), as well as the dBP control group ($I^2=0.47$). The Egger's regression test was performed, which was only significant for dBP in the IET trials ($P= 0.04$) (Figure S1).

High intensity interval training

HIIT significantly reduced sBP and dBP by a mean difference of 2.86mmHg (95% CI= 1.11 to 4.62, $p=0.0014$) and 2.48 mmHg (95% CI= 1.49 to 3.48, $p<0.001$) respectively, with no significant sBP change observed in the control group, but a statistically significant increase in the control group dBP (MD: -0.50, 95% CI= -0.961 to -0.041, $p=0.03$). mBP also significantly decreased following HIIT by 3.15 mmHg (95% CI= 0.96 to 5.34, $p=0.0049$) with no significant change in the control group. There were significant differences between groups for dBP ($p<0.001$) and mBP ($p=0.038$), with a (MD: 1.03, 95% CI= -0.170 to 2.224, $p=0.09$). HIIT also

had a significant effect on RHR by a mean difference of 3.17 bpm (95% CI= 2.49 to 3.85, $p<0.001$) with no changes in the control group and a significant between-groups effect.

There was significant heterogeneity between studies in the HIIT sBP ($I^2=52\%$), dBP ($I^2=34\%$) and mBP arms ($I^2=47\%$), with no significant heterogeneity in the control group. Eggers test (1997) for publication bias was significant for asymmetry in sBP in the HIIT trials ($P =0.02$) (Figure S2).

Isometric exercise training vs high intensity interval training

There was a significant treatment difference between IET and HIIT trials for all variables. IET produced significantly greater reductions in resting BP compared to HIIT in sBP (MD: 5.29, 95% CI = 3.97 to 6.61, $Q = 17.1$, $p<0.01$, $I^2 = 86\%$; Figure 2), dBP (MD: 3.25, 95% CI = 2.53 to 3.96, $Q = 4.71$, $p=0.03$, $I^2 = 70\%$; Figure 3) and mBP (MD: 4.63, 95% CI = 3.09 to 6.17, $Q = 4.54$, $P=0.03$, $I^2 = 83\%$). However, HIIT reduced RHR to a significantly greater extent than IET (MD: 2.45, 95% CI = 1.78 to 3.11, $Q=6.01$, $p=0.01$, $I^2 =22\%$).

Meta-regression

There was no statistical significance for the moderator variables: hypertension diagnosis, medication status and intervention duration. As such, TESTEX score was the only statistically significant moderator, which explained a significant proportion of the dBP inter-study variance in the HIIT trials only ($R^2 = 75.41$, $p=0.014$, 95% CI= 0.137 to 1.205) (Figure S3).

Discussion

The aim of this study was to perform a meta-analytic comparison of randomised controlled trials to directly compare the efficacy of IET versus HIIT in the management of BP. While the findings of our analysis support both IET and HIIT as efficacious modalities in reducing resting sBP, dBP and mBP, this novel comparison reported a statistically significant between-treatment difference, with reductions of a greater magnitude in all BP measures following IET compared to HIIT. However, resting heart rate (RHR) was reduced to a significantly greater extent following HIIT compared to IET.

IET demonstrated significant sBP, dBP and mBP reductions of -8.5, -4.07 and -6.46 mmHg respectively. Importantly, these reductions are clinically significant and of a similar, or greater magnitude to the reductions commonly observed following standard anti-hypertensive pharmacotherapy [21]. Although the lower limit confidence intervals were below what is generally accepted as clinically significant, HIIT produced significant reductions of -2.86, -2.48 and -3.15 in sBP, dBP and mBP respectively. While not to the same magnitude as IET, these HIIT reductions are certainly comparable to the effects typically seen following traditional moderate intensity aerobic exercise [22] which is currently recognised as the recommended modality for the management of BP in international guidelines [23]. Indeed, a previous meta-analysis [22] reported reductions in resting sBP and dBP by -3.84 and -2.58 mmHg following traditional aerobic exercise, thus demonstrating the comparative efficacy of these modes and highlighting IET as unequivocally superior. Ultimately, the presented anti-hypertensive effects of both IET and HIIT are associated with a reduced risk of cardiovascular disease and all-cause mortality, therefore providing imperative clinical implications regarding the optimal control of BP [24,25].

The mechanistic underpinnings behind the anti-hypertensive effects of IET and HIIT are highly complex and still largely unknown, but are likely to share some homogenous physiological processes [26,27]. Despite the significant reductions in RHR observed in our analysis, both interventions have often been shown to reduce BP with only small or no changes in CO [8,10]. This suggests that BP is reduced following either intervention primarily via changes in total peripheral resistance. As such, both modes have been linked with enhanced shear stress as a mechanical stimulus to facilitate increases in endothelial intracellular calcium via potassium channel activation, ultimately promoting endothelial nitric

oxide synthase and thus nitric oxide bioavailability [28]. While this mechanism is conceptually consistent across both modes, the stimulating process of flow mediated dilation (FMD) differs between IET and HIIT within an acute setting [27,29,30], which although speculative, may be implicit in determining the observed differences in magnitude of BP reduction. Specifically, IET acutely produces such a mechanical response via contraction-induced occlusion of the relevant vasculature, thus causing reactive hyperaemia on relaxation to produce such mechanism, which has been demonstrated to translate into chronic FMD changes adjacent to significant improvements in BP [31]. Conversely, HIIT simply invokes standard exercise hyperaemia, and despite the considerable stress placed on the relevant skeletal muscles with such workload intensity, current evidence indicates that the shear rate following HIIT is no greater than that of traditional aerobic exercise, thus potentially limiting the anti-hypertensive effects [28,32]. However, the mechanistic rationale behind such differences in BP reducing effects between both modes is not understood and requires future investigation.

Interestingly, we found no significant differences for any of the measured parameters between upper (handgrip) and lower (wall-squat and leg-extension) body IET modes. While handgrip remains the traditionally investigated IET mode, more recent data has demonstrated lower-body IET to achieve reductions to a much larger magnitude [8]. Such differences have been mechanistically attributed to the larger muscle mass involvement in lower body IET, generating a greater occlusion-response stimulus and thus enhancing the metabolic and autonomic regulatory adaptations [8,29]. To what extent the lack of differences observed in the present analysis is owing to the disparity in published trials between IET modes is unknown, with future lower body IET research warranted before definitive conclusions on the optimal application of IET can be made. Furthermore, we also found no significant effects of medication and hypertension status on BP or RHR changes following IET or HIIT. Similarly, this finding may also be somewhat attributable to methodological limitations, with many studies failing to report exact participant characteristic data on medication or hypertension status, as well as the role of different medications and varying severities of hypertension increasing the complexity of this analysis.

Separately, both modes also show significant reductions in RHR with a significantly greater magnitude of change following HIIT compared to IET, perhaps indicating a more prominent role of RHR in the observed BP changes following HIIT than IET. Given the strong

association between RHR and cardiovascular and all-cause mortality [33], these findings may be of clinical importance. Both IET and HIIT have been shown to produce significant improvements in cardiac autonomic regulation and baroreceptor sensitivity [8,10], which as well as mechanistically contributing to the observed BP changes, are likely to be implicit in these RHR reductions. Additionally, previous investigations from our lab in HIIT [10] and acute IET [34] have demonstrated significant improvements in cardiac function and mechanics following both interventions, providing potential mechanistic implications for such RHR changes. In speculating the differences in the magnitude of RHR change, it is well-established that, despite being an anaerobic modality in nature, HIIT may induce significant peripheral and haematological adaptations that are generally associated with traditional aerobic exercise, such as increased skeletal muscle mitochondrial content [35,36], capillary density [37] and plasma and blood volume [38], thus contributing to enhanced arteriovenous oxygen difference and stroke volume, resulting in vagally-mediated reductions in resting RHR to a greater extent [27]. Conversely, such physiological changes are not generally associated with IET.

Aside from RHR, these physiological adaptations raise an important point in distinguishing the appropriate application of either modality. In particular, these HIIT-induced adaptations may directly contribute to functional capacity and therefore have significant implications for clinical prognosis as one of the strongest predictors of outcome [39]. Thus, it may be speculated that, while IET is superior in the management of resting BP, especially in hypertensive populations, HIIT may provide wider physiological advantages and greater clinical implications in individuals without hypertension, making it the preferred modality for the maintenance of general health. Therefore, since both modes may have different physiological advantages, future research should explore the application of both IET and HIIT in combination versus single-mode exercise training.

Limitations

As a primary limitation, we found significant heterogeneity for all BP outcomes in both interventions. Attributing this variance to methodological differences, our meta-regression accounted for hypertension diagnosis, medication status and intervention duration, but found no statistical significance for these moderators. Risk of bias TESTEX score explained a significant proportion of the HIIT DBP inter-study variance only. Nonetheless, random-effects models were applied to account for such heterogeneity. The Eggers plots showed

significant publication bias for sBP in the IET trials and dBP in the HIIT trials, suggesting studies that found no significant reductions in BP following either intervention may not have been published. Furthermore, the participants and investigators would have been aware of group allocation, as assessed in the TEXTEX scale. Thus, future IET and HIIT research should attempt to employ a sham-design methodology to blind the participants to their group allocation. It should also be considered that the present within-group analyses are subject to various biases, such as regression to the mean. There were also search limitations as we applied an English language filter, which carries the risk of introducing selection bias, and we did not search Scopus or Web of Science databases. Finally, this analysis did not compare IET or HIIT against MICT, which remains the most commonly applied mode, thus limiting the wider interpretations of the present results.

Clinical Implications

Hypertension remains the leading risk factor for cardiovascular disease and all-cause mortality worldwide [1]. Given the widely documented limitations that come with medication usage, it is crucial that effective non-pharmacological / lifestyle treatment options are established. IET and HIIT are short duration, highly time-efficient modes which have been shown to produce clinically significant reductions in resting BP at a magnitude superior to that of traditional exercise training modalities [7,9]. This study demonstrates the efficacy of both interventions in the management of BP, highlighting IET as the superior mode in reducing BP, while HIIT remains more efficacious in reducing RHR. Specifically, at a magnitude of 8.5mmHg sBP and 4mmHg dBP, IET may produce reductions in BP to a greater extent than that typically achieved with standard dose anti-hypertensive monotherapy [40]. While less effective for BP, HIIT may induce wider cardiovascular adaptations, making it the preferred mode in the general maintenance of health.

Conclusion

Both IET and HIIT are efficacious modes in the management of resting BP. IET appears the superior mode, producing a greater magnitude of BP reduction when compared to HIIT. However, HIIT remains a more effective mode in reducing RHR. Future research, ideally employing a sham-design protocol, is needed to establish the mechanistic underpinnings behind such differences in BP reducing effects and to explore the efficacy of both modes applied in combination.

Competing interests: None

Contributorship: JE, AD and JO'D contributed to the conception and design of the study. JE and AD contributed equally to this work. JE, AD and JO'D contributed to the development of the search strategy. JE and AD conducted the systematic review. JE, AD and JO'D completed the acquisition of data. JE, AD, JD and JO'D performed the data analysis. All authors assisted with the interpretation. JE and JO'D were the principal writers of the manuscript. All authors contributed to the drafting and revision of the final article. All authors approved the final submitted version of the manuscript.

Acknowledgments: The authors would like to thank Michelle Crowther who is Canterbury Christ Church Universities learning and research librarian. Michelle assisted in developing the search strategy for this meta-analysis.

Fundng, Grant and Awards: None

Ethical Approval: Not applicable

Data Sharing Agreement: Not applicable

Reference List

- 1 Lim SS, Vos T, Flaxman AD, *et al.* A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;**380**:2224–60. doi:10.1016/S0140-6736(12)61766-8
- 2 Gebreyohannes EA, Bhagavathula AS, Abebe TB, *et al.* Adverse effects and non-adherence to antihypertensive medications in University of Gondar Comprehensive Specialized Hospital. *Clin Hypertens* 2019;**25**:1–9. doi:10.1186/s40885-018-0104-6
- 3 Wang G, Grosse SD, Schooley MW. Conducting Research on the Economics of Hypertension to Improve Cardiovascular Health. *Am. J. Prev. Med.* 2017;**53**:S115–7. doi:10.1016/j.amepre.2017.08.005
- 4 Vrijens B, Vincze G, Kristanto P, *et al.* Adherence to prescribed antihypertensive drug treatments: Longitudinal study of electronically compiled dosing histories. *BMJ* 2008;**336**:1114–7. doi:10.1136/bmj.39553.670231.25
- 5 Williamson W, Foster C, Reid H, *et al.* Will Exercise Advice Be Sufficient for Treatment of Young Adults With Prehypertension and Hypertension? A Systematic Review and Meta-Analysis. *Hypertension*. 2016;**68**:78–87. doi:10.1161/HYPERTENSIONAHA.116.07431
- 6 Guthold R, Stevens GA, Riley LM, *et al.* Worldwide trends in insufficient physical activity from 2001 to 2016: a pooled analysis of 358 population-based surveys with 1·9 million participants. *Lancet Glob Heal* 2018;**6**:e1077–86. doi:10.1016/S2214-109X(18)30357-7
- 7 Inder JD, Carlson DJ, Dieberg G, *et al.* Isometric exercise training for blood pressure management: A systematic review and meta-analysis to optimize benefit. *Hypertens Res* 2016;**39**:89–94. doi:10.1038/hr.2015.111
- 8 Taylor KA, Wiles JD, Coleman DA, *et al.* Neurohumoral and ambulatory haemodynamic adaptations following isometric exercise training in unmedicated hypertensive patients. *J Hypertens* 2019;**37**:827–36. doi:10.1097/HJH.0000000000001922
- 9 Edwards JJ, Taylor KA, Cottam C, *et al.* Ambulatory blood pressure adaptations to high-intensity interval training: a randomized controlled study. *J Hypertens* 2021;**39**:341–8. doi:10.1097/HJH.0000000000002630
- 10 O’Driscoll JM, Wright SM, Taylor KA, *et al.* Cardiac autonomic and left ventricular

- mechanics following high intensity interval training: A randomized crossover controlled study. *J Appl Physiol* 2018;**125**:1030–40.
doi:10.1152/jappphysiol.00056.2018
- 11 Herazo-Beltrán Y, Pinillos Y, Vidarte J, *et al.* Predictors of perceived barriers to physical activity in the general adult population: a cross-sectional study. *Brazilian J Phys Ther* 2017;**21**:44–50. doi:10.1016/j.bjpt.2016.04.003
 - 12 Costa EC, Hay JL, Kehler DS, *et al.* Effects of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training On Blood Pressure in Adults with Pre- to Established Hypertension: A Systematic Review and Meta-Analysis of Randomized Trials. *Sport. Med.* 2018;**48**:2127–42. doi:10.1007/s40279-018-0944-y
 - 13 Cornelissen VA, Smart NA. Exercise training for blood pressure: a systematic review and meta-analysis. *J. Am. Heart Assoc.* 2013;**2**. doi:10.1161/JAHA.112.004473
 - 14 López-Valenciano A, Ruiz-Pérez I, Ayala F, *et al.* Updated systematic review and meta-analysis on the role of isometric resistance training for resting blood pressure management in adults. *J. Hypertens.* 2019;**37**:1320–33.
doi:10.1097/HJH.0000000000002022
 - 15 Hansen D, Dendale P, Coninx K, *et al.* The European Association of Preventive Cardiology Exercise Prescription in Everyday Practice and Rehabilitative Training (EXPERT) tool: A digital training and decision support system for optimized exercise prescription in cardiovascular disease. Concept, definitions and construction methodology. *Eur J Prev Cardiol* 2017;**24**:1017–31. doi:10.1177/2047487317702042
 - 16 Smart NA, Waldron M, Ismail H, *et al.* Validation of a new tool for the assessment of study quality and reporting in exercise training studies: TESTEX. *Int J Evid Based Healthc* 2015;**13**:9–18. doi:10.1097/XEB.0000000000000020
 - 17 Balduzzi S, Rucker G, Schwarzer G. How to perform a meta-analysis with R: A practical tutorial. *Evid Based Ment Health* 2019;**22**:153–60. doi:10.1136/ebmental-2019-300117
 - 18 Fu R, Holmer HK. Change score or follow-up score? Choice of mean difference estimates could impact meta-analysis conclusions. *J Clin Epidemiol* 2016;**76**:108–17.
doi:10.1016/j.jclinepi.2016.01.034
 - 19 Rothman, Kenneth J; Greenland, Sander; Lash TL. *Modern Epidemiology*. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins 2008.
 - 20 Harrer M, Cuijpers P, Furukawa T, *et al.* dmetar: Companion R Package For The Guide ‘Doing Meta-Analysis in R’. R package version 0.0.9000.

- 2019.<http://dmetar.proctclab.org/>
- 21 Paz MA, De-La-Sierra A, Sáez M, *et al.* Treatment efficacy of anti-hypertensive drugs in monotherapy or combination: ATOM systematic review and meta-analysis of randomized clinical trials according to PRISMA statement. *Med. (United States)*. 2016;**95**. doi:10.1097/MD.0000000000004071
- 22 Whelton SP, Chin A, Xin X, *et al.* Effect of aerobic exercise on blood pressure: A meta-analysis of randomized, controlled trials. *Ann Intern Med* 2002;**136**:493–503. doi:10.7326/0003-4819-136-7-200204020-00006
- 23 World Health Organization. Global recommendations on physical activity for health. 2010.
<http://medcontent.metapress.com/index/A65RM03P4874243N.pdf%5Cnhttp://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Global+Recomendations+on+physical+activity+for+health#0> (accessed 3 Apr 2020).
- 24 Brunström M, Carlberg B. Association of blood pressure lowering with mortality and cardiovascular disease across blood pressure levels a systematic review and meta-analysis. *JAMA Intern. Med.* 2018;**178**:28–36. doi:10.1001/jamainternmed.2017.6015
- 25 Ettehad D, Emdin CA, Kiran A, *et al.* Blood pressure lowering for prevention of cardiovascular disease and death: A systematic review and meta-analysis. *Lancet* 2016;**387**:957–67. doi:10.1016/S0140-6736(15)01225-8
- 26 Millar PJ, McGowan CL, Cornelissen VA, *et al.* Evidence for the role of isometric exercise training in reducing blood pressure: Potential mechanisms and future directions. *Sport. Med.* 2014;**44**:345–56. doi:10.1007/s40279-013-0118-x
- 27 MacInnis MJ, Gibala MJ. Physiological adaptations to interval training and the role of exercise intensity. *J Physiol* 2017;**595**:2915–30. doi:10.1113/JP273196
- 28 Ramos JS, Dalleck LC, Tjonna AE, *et al.* The Impact of High-Intensity Interval Training Versus Moderate-Intensity Continuous Training on Vascular Function: a Systematic Review and Meta-Analysis. *Sport. Med.* 2015;**45**:679–92. doi:10.1007/s40279-015-0321-z
- 29 Wiles JD, Goldring N, Coleman D. Home-based isometric exercise training induced reductions resting blood pressure. *Eur J Appl Physiol* 2017;**117**:83–93. doi:10.1007/s00421-016-3501-0
- 30 Taylor KA, Wiles JD, Coleman DD, *et al.* Continuous cardiac autonomic and hemodynamic responses to isometric exercise. *Med Sci Sports Exerc* 2017;**49**:1511–9. doi:10.1249/MSS.0000000000001271

- 31 Badrov M, Freeman S, Zokvic M, *et al.* Isometric exercise training lowers resting blood pressure and improves local brachial artery flow-mediated dilation equally in men and women. *Eur J Appl Physiol* 2016;**116**:1289–96.
<http://search.ebscohost.com/login.aspx?direct=true&db=s3h&AN=116194061&site=ehost-live>
- 32 Currie KD, Dubberley JB, McKelvie RS, *et al.* Low-volume, high-intensity interval training in patients with CAD. *Med Sci Sports Exerc* 2013;**45**:1436–42.
doi:10.1249/MSS.0b013e31828bbbd4
- 33 Zhang D, Shen X, Qi X. Resting heart rate and all-cause and cardiovascular mortality in the general population: A meta-analysis. *CMAJ* 2016;**188**:E53–63.
doi:10.1503/cmaj.150535
- 34 O’Driscoll JM, Taylor KA, Wiles JD, *et al.* Acute cardiac functional and mechanical responses to isometric exercise in prehypertensive males. *Physiol Rep* 2017;**5**.
doi:10.14814/phy2.13236
- 35 Talanian JL, Galloway SDR, Heigenhauser GJF, *et al.* Two weeks of high-intensity aerobic interval training increases the capacity for fat oxidation during exercise in women. *J Appl Physiol* 2007;**102**:1439–47. doi:10.1152/jappphysiol.01098.2006
- 36 MacInnis MJ, Zacharewicz E, Martin BJ, *et al.* Superior mitochondrial adaptations in human skeletal muscle after interval compared to continuous single-leg cycling matched for total work. *J Physiol* 2017;**595**:2955–68. doi:10.1113/JP272570
- 37 Cocks M, Shaw CS, Shepherd SO, *et al.* Sprint interval and endurance training are equally effective in increasing muscle microvascular density and eNOS content in sedentary males. *J Physiol* 2013;**591**:641–56. doi:10.1113/jphysiol.2012.239566
- 38 Graham MJ, Lucas SJE, Francois ME, *et al.* Low-Volume Intense Exercise Elicits Post-exercise Hypotension and Subsequent Hypervolemia, Irrespective of Which Limbs Are Exercised. *Front Physiol* 2016;**7**:199. doi:10.3389/fphys.2016.00199
- 39 Kodama S, Saito K, Tanaka S, *et al.* Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: A meta-analysis. *JAMA - J. Am. Med. Assoc.* 2009;**301**:2024–35.
doi:10.1001/jama.2009.681
- 40 Law MR, Morris JK, Wald NJ. Use of blood pressure lowering drugs in the prevention of cardiovascular disease: Meta-analysis of 147 randomised trials in the context of expectations from prospective epidemiological studies. *BMJ* 2009;**338**:1245.
doi:10.1136/bmj.b1665

Table 1. IET study characteristics.

Study	Country	Duration (weeks)	Participants	Hypertension	Medication	Withdrawal (N. of participants)	Training Frequency	Exercise Mode	Exercise Training Characteristics	TESTEX score
Baddeley- White et al (2019)	UK	4	n=23 (13 Male &10 Female) 23 <50 years	NTN	Medicated= 0	None	3 x per week	Isoball rugby handgrip/ zona plus handgrip	4 x 2 min, 1 min rest interval, 30%MVC, (n=7 isoball, n=8 zona, n=8 control).	7
Badrov et al (2013a)	Canada	8	n=36 (36 Female) 36 <50 years	NTN	Medicated= 0	IET= 1 Control= 3	3/5 x per week	Handgrip	4 x 2 min, 4 min rest intervals, 30% MVC, (n=12 3x per week, n=11 5 x per week, n=9 control).	10
Badrov et al (2013b)	Canada	10	n=24 (13 Male & 11 Female) 24 >50 years	HTN	Medicated= 0	None	3 x per week	Handgrip	4 x 2 min bilateral, 1 min rest interval, 30% MVC, (n=12 IET, n=12 control).	8
Baross et al (2012)	UK	8	n=30 (30 Male) 30 >50 years	Mixed NTN/HTN	Medicated= 0	None	3 x per week	Leg extension (Bilateral)	4 x 2 min, 2 min rest intervals, 14%MVC, (n=10 at 85%HRpeak, n= 10	7

Author (Year)	Country	Study ID	Participants (n)	Condition	Medicated	Control	Frequency	Exercise	Protocol	Outcome
Baross et al (2013)	UK	8	n=20 (20 Male) 20 >50 years	NTN	Medicated = 0	None	3 x per week	Leg extension (Bilateral)	4 x 2 min, 2 min rest intervals, (85%HRpeak n=10 exercise group, n=10 control).	7
Cahu Rodrigues et al (2019)	Brazil	12	n=72 (67% Female) (age unknown)	HTN	Medicated= 72	IET=31 Control=8	3 x per week	Handgrip	4 x 2 min, 1 min rest interval, (n=17 30%MVC, n=16 control).	10
Carlson et al (2016)	Australia	8	n=40 (15 Male & 25 Female) 40 >50 years	HTN	Medicated= 26	IET= 2	3 x per week	Handgrip	4 x 2 min, 1 min rest intervals, (n= 18 at 30%MVC, n=20 5%MVC exercise control).	13
Correia et al (2020)	Brazil	8	n=102 (sex unknown)	Pre-HTN	Medicated= 0	IET=21 Control= 2	3 x per week	Handgrip	4 x 2 min, 4 min rest intervals, (n=29 30%MVC, n=50 control).	9

			102 >50 years							
Farah et al (2018)	USA	12	n=72 (75% Female) 48 >50 years	HTN	Medicated= 72	Home IET= 6 Supervised IET= 10 Control= 8	3 x per week	Handgrip	4 x 2 min, 1 min rest intervals, (n=13 at 30%MVC home based, n=14 30%MVC supervised, n=16 non exercise control).	9
Gordon et al (2018)	USA	12	n=22 (sex unknown) 22 <50 years	HTN	Medicated= 20	None	3 x per week	Handgrip	4 x 2 min, 1 min rest intervals, (n=5 30%MVC home based, n=8 30%MVC lab based, n=9 control).	8
Miller et al (2013)	Canada	8	n=23 (18 Male & 5 Female) 23 >50 years	HTN	Medicated= 23	None	3 x per week	Handgrip	4 x 2 min, 1 min rest intervals, (n=13 30%MVC, n=10 control).	7

Okamoto et al (2020)	Japan	8	n=22 (9 Male & 13 Female) 22 >50 years	Pre-HTN	Medicated=0	None	3 x per week	Handgrip	4 x 2 min, 1 min rest interval. (n=11 30% MVC, n=11 control).	10
Punia et al (2019)	India	8	n=40 (Male & Female) 40 <50 years	HTN	Mixed Medicated & Non-Medicated	None	3 x per week	Handgrip	4 x 2 min, 4 min rest intervals, (n=20 30%MVC, n=20 non-exercise control group).	10
Taylor et al (2003)	Canada	10	n=17 (10 Male & 7 Female) 17 <50 years	HTN	Medicated=75%	None	3 x per week	Handgrip	4 x 2 min, 1 min rest intervals, (n=9 30%MVC, n=8 non-exercise control group).	7
Taylor et al (2018)	UK	4	n=48 (Male & Female) (age unknown)	HTN	Medicated=0	None	3 x per week	Wall squat	4 x 2 min, 2 min rest intervals, (n=24 95% HRpeak, n=24 non exercise control).	7
Wiles et al (2009)	UK	8	n=33 (33 Males) 33 <50 years	NTN	Medicated=0	None	3 x per week	Leg extension (Bilateral)	4 x 2 min, 2 min rest intervals, (n=11 HI-95%HRpeak, n=11	8

Author (Year)	Country	Interventions	n	NTN	Medicated	None	Frequency	Exercise	Control	Score
Wiles et al (2016)	UK	4	n=28 (28 Male) 28 <50 years	NTN =28	Medicated= 0	None	3 x per week	Wall squat	LO-75%HRpeak, n=11 control group). 4 x 2 min, 1 min rest interval, (n=14 95%HRpeak, n=14 non-exercise control).	7
Yamagata et al (2020)	Japan	8	n=20 (sex and age unknown)	NTN= 20	Medicated= 0	None	3 x per week	Handgrip	4 x 2 min, 3 min rest intervals. (n=10 25% MVC Handgrip, n=10 control group).	9

Abbreviations; IET, isometric exercise training, HI, high, HR, heart rate, LO, low, MVC, maximum voluntary contraction, HTN, hypertensive, NTN, normotensive, pre-HTN, pre-hypertensive; RPE, rate of perceived exertion.

Table 2. HIIT study characteristics.

Study	Country	Duration (weeks)	Participants	Hypertension	Medication	Withdrawal (N. of participants)	Training Frequency	Exercise Mode	Exercise Training Characteristics	TESTEX score
Amaro-Gahete et al (2019)	Spain	12	n=89 (32 Male and 37 Female) 69 >50 years	Mixed HTN/NTN	Medication not reported	PA= 4 Control= 5 HIIT=5 HIIT+EMS=4	2 x per week	Sloped treadmill and weighted exercises	Walking on an incline, n=17 PA (150min p/w), n=18 HIIT-LI (40-60min p/w at 95% VO ² max) and HIIT-SI (16 weighted exercise 6-9 RPE), n=19 HIIT+EMS, (HIIT with electromyostimulation), n=17 non-exercise control.	11
Astorino et al (2012)	USA	2	n=29 (16 Male and 13 Female) 29 <50 years	NTN	Medication not reported	None	3 x per week	Cycling	4 x 30 sec with 5 min rest intervals, n=20 HIIT 7.5% BW all out max, n=9 control.	6
Boutcher et al (2019)	Australia	8	n=40 (40 Female) 40 >50years	NTN	Medication not reported	None	3 x per week	Cycling	8 sec SIT intervals with 12 sec rest intervals for 20 min, n=20 SIT at 80-85%HRpeak, n=20 non-exercise control.	9
Cassidy et al (2019)	UK	12	n=22 (17 Male and 5 Female)	NTN	Medicated n= 8	None	3 x per week	Cycling	5 x 2 min increasing by 10 sec a week up to 3 min 50 sec by week 12 with 90 sec	10

			11 >50 years						passive recovery and 60 sec band exercises, n=11 HIIT 16-17 RPE and 80 revolution cadence, n=11 control.	
Dall et al (2014)	Denmark	12	n=17 (12 Male 5 Female) 17 >18 years	HTN= 81%	Medicated n= 94%	Control= 1	3 x per week	Cycling	4, 2, 1 min intervals for 16 mins, 2 min rest intervals (32 min session), n=8 >80% VO ² peak, n=8 control 45 min continuous aerobic 60% VO ² max.	10
Edwards et al (2020)	UK	4	n=41 (20 Male 21 Female) Age not stated	NTN	Medicated n= 0	None	3 x per week	Cycling	3 x 30 sec intervals, 2 min rest intervals, n=21 at 7.5%bw, n=20 control group.	8
Gerosa-Neto et al (2019)	Brazil	6	n=46 (46 Male) 46 <50 years	Mixed HTN/NTN	Medication not reported	HIIT= 5 MICT= 5 Control= 18	3 x per week	Treadmill	10 x 1 min intervals, 1 min rest intervals, n=18 HIIT at 100% VO ² max, n=18 MICT.	7
Ghardashi afousi et al (2018)	Iran	12	n=75 (25 Males, 27 Female) 52 >50 years	Mixed HTN/NTN	Medicated n= ~50%	HIIT= 7 MICT= 8 Control= 8	3 x per week	Cycling	12 x 1.5 min intervals, 2 min rest intervals, n=18 at 85%-90% HRmax and 2 min at 55%-60% HRmax, n=17 42 min at 70% HRmax, n=18 control.	8

Hanssen et al (2017)	Switzerland	12	n=48 (7 Males, 30 Female) 48 <50 years	Mixed HTN/NTN	Medication not reported	HIIT= 3 MICT= 4 Control= 4	2 x per week	Treadmill running	4 x 4 min intervals, 3 min rest intervals, n=13 at 90-95% HRmax, n=12 45 min 70% HRmax, n=12 control.	7
Hallsworth et al (2015)	UK	12	n=28 (all same sex not stated if Male or Female) 23 >50 years	Mixed HTN/NTN	Medication not reported	HIIT= 2 Control= 3	3 x per week	Cycling	5 x 2 min intervals, 3 min rest intervals, increase 10 sec weekly, n=12 HIIT at 16-17 RPE, n= 11 control.	8
Izadi et al (2018)	Iran	6	n=44 (17 Male 13 Female) 44 >50 years	NTN	Medicated n= 44	HIIT= 7 Control= 7	3 x per week	Cycling	10 x 1.5 min intervals, 2 min rest interval (35 min session), n=15 HIIT 90-95% HRR, n=15 control.	8
Jabbour et al (2016)	Canada	6	n=24 (12 Male 12 Female) 24 <50 years	Mixed HTN/NTN	Medication not reported	None	3 x per week	Cycling	6 x 6 sec intervals, 2 min rest intervals, n=12 HIIT at maximal output, n=12 control group.	7
Jung et al (2015)	Canada	2	n=32 (5 Male 27 Female) 32 >50 years	Mixed HTN/NTN	Medication not reported	HIIT= 5 Control= 1	5 x per week	Cycling, treadmill, cross-trainer	4 x 1 min intervals, increasing up to 10 x 1 min by day 10, 1 min rest intervals, n=15 HIIT at	10

									90% HRpeak, n=17 control at 60% HRpeak.	
Lee et al (2019)	USA	8	n=30 (30 Female) 30 >50 years	Mixed HTN/NTN	Medication not reported	None	3 x per week	Cycling	7 x 1 min intervals, 2 min rest intervals (20 min session), n=15 HIIT at 90% peak power output, n=15 control group.	8
May et al (2018)	USA	4	n=90 (82% Female) 90 <50 years	NTN	Medicated n= 0	None	3 x per week	Cycling	10 x 1 min intervals, 1 min rest interval, n=30 HIIT at 90% HRmax, n=30 HRVCB training, n=30 control.	8
O'Driscoll et al (2018)	UK	2	n=44 (44 Female) 44 <50 years	NTN	Medicated n= 0	HIIT= 2 Control= 2	3 x per week	Cycling	3 x 30 sec Wingate intervals, 2 min rest intervals, n=20 HIIT 7.5%bw, n=20 control.	8
Reljic et al (2020)	Germany	12	n=65 (29 Male 36 Female) 65 <50 years	Mixed HTN/NTN	Medicated= mixed, data not reported	n=16	2 x per week	Cycling	5 x 1 min intervals, 1 min rest intervals, n=36 at 80-95%HRmax, n=29 control.	13
Rognmo et al (2004)	Norway	10	n=21 (14 Male 3 Female) 21 >50 years	Mixed HTN/NTN	Medicated= yes, data not reported	HIIT= 3 Control= 1	3 x per week	Incline treadmill	4 x 4 min intervals, 3 min rest intervals, n=8 at 90% HRpeak, n=9 control.	13

Sandstad et al (2015)	Norway	10	n=36 (gender not stated) 36 <50 years	NTN	Medication not reported	HIIT= 6 Control= 3	2 x per week	Cycling	4 min intervals, 3 min rest intervals for 35 min, n= 12 HIIT at 95% HRpeak, n=15 control group.	8
Shepherd et al (2015)	UK	10	n=90 (30 Male 60 Female) 90 <50 years	Mixed HTN/NTN	Medication not reported	None	3 x per week	Cycling	15-60 sec intervals, 45-120 sec rest intervals (8-25 min session), n=46 at > 90% HRmax, n=44 MICT at 70% HRpeak.	11

Abbreviations: HIIT, high intensity interval training; HI, high intensity; LI, low intensity; HR, heart rate; MVC, maximum voluntary contraction; HTN, hypertensive; NTN, normotensive; pre-HTN, pre-hypertensive; RPE, rate of perceived exertion.

Figure Legend:

Figure 1: PRISMA systematic review and meta-analysis flowchart.

Figure 2: Random-effects meta-analysis of the effects of IET versus HIIT in the reduction of sBP.

Figure 3: Random-effects meta-analysis of the effects of IET versus HIIT in the reduction of dBP

What is already known

- Anti-hypertensive pharmacotherapy carries limitations and the current exercise guidelines for the control of resting blood pressure are limited by poor compliance.
- Isometric exercise training and high-intensity interval training have been shown to improve markers of cardiovascular health.
- Both modalities have promising future clinical utility in the management of resting blood pressure.

What are the new findings

- Compared to high intensity interval training, isometric exercise training is the superior modality in the management of resting blood pressure, with reductions similar to, or greater than that of anti-hypertensive medication.
- While still efficacious in reducing resting blood pressure, high-intensity interval training may achieve wider physiological benefits, with greater reductions in resting heart rate.
- As such, the clinical implementation of isometric exercise training for blood pressure control should be considered, with high-intensity interval training likely preferred for the general maintenance of optimal health.