

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Cerebroplacental ratio at 35-37 weeks' gestation in the prediction of adverse perinatal outcome

Ranjit AKOLEKAR, M.D.,<sup>1,2</sup> Anca CIOBANU, M.D.,<sup>3</sup> Emilie ZINGLER, M.D.<sup>3</sup> Argyro SYNGELAKI, Ph.D.,<sup>3</sup> Kypros H. NICOLAIDES, M.D.<sup>3</sup>

1. Fetal Medicine Unit, Medway Maritime Hospital, Gillingham, UK
2. Institute of Medical Sciences, Canterbury Christ Church University, Chatham, UK
3. Fetal Medicine Research Institute, King's College Hospital, London, UK

**Short title:** Third trimester screening for SGA

**Key words:** Third trimester screening, Small for gestational age, Umbilical artery Doppler, Middle cerebral artery Doppler, Cerebroplacental ratio; Adverse perinatal outcome.

### Correspondence:

Professor KH Nicolaides,  
Fetal Medicine Research Institute,  
King's College Hospital,  
16-20 Windsor Walk,  
Denmark Hill, London SE5 8BB  
Telephone: +442032998256  
Fax: +442077339534  
email: [kypros@fetalmedicine.com](mailto:kypros@fetalmedicine.com)

**Acknowledgement:** The study was supported by grants from the Fetal Medicine Foundation (Charity No: 1037116). This body had no involvement in the study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication.

## CONDENSATION

Routine assessment of cerebroplacental ratio at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation provides poor prediction of adverse perinatal outcome.

## AT A GLANCE

- A. To investigate the performance of screening for adverse perinatal outcome by the cerebroplacental ratio (CPR) measured routinely at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation.
- B. In a prospective observational study in 47,211 women with singleton pregnancies undergoing routine ultrasound examination at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation, low CPR was associated with increased risk of adverse perinatal outcome, presence of surrogate markers of perinatal hypoxia, cesarean section for presumed fetal distress in labor and birth of neonates with birthweight <3<sup>rd</sup> percentile. However, the performance of low CPR in the prediction of each adverse outcome was poor, with detection rates of 13–26% and false positive rate of about 10%.
- C. In pregnancies undergoing routine antenatal assessment at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation measurement of CPR provides poor prediction of adverse perinatal outcome in both small and appropriate for gestational age fetuses. Consequently, there is no justification in a shift of the focus of prenatal care from identification of pregnancies with low estimated fetal weight to that of pregnancies with low CPR.

## ABSTRACT

**Background:** Third trimester studies in selected high-risk pregnancies have reported that low cerebroplacental ratio (CPR), due to high pulsatility index (PI) in the umbilical artery (UA), and or decreased PI in the fetal middle cerebral artery (MCA), is associated with increased risk of adverse perinatal outcomes.

**Objective:** To investigate the predictive performance of screening for adverse perinatal outcome by the cerebroplacental ratio (CPR) measured routinely at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation.

**Methods:** This was a prospective observational study in 47,211 women with singleton pregnancies undergoing routine ultrasound examination at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation, including measurement of UA-PI and MCA-PI. The measured UA-PI and MCA-PI and their ratio were converted to multiples of the median (MoM) after adjustment for gestational age. Multivariable logistic regression analysis was used to determine whether CPR improved the prediction of adverse perinatal outcome that was provided by maternal characteristics, medical history and obstetric factors. The following outcome measures were considered: first, adverse perinatal outcome consisting of stillbirth, neonatal death or hypoxic ischemic encephalopathy grades 2 and 3, second, presence of surrogate markers of perinatal hypoxia consisting of umbilical arterial or venous cord blood pH  $\leq 7$  and  $\leq 7.1$ , respectively, 5-minute Apgar score  $< 7$ , or admission to the neonatal intensive care unit for  $> 24$  hours, third, cesarean section for presumed fetal distress in labor, and fourth, neonatal birthweight  $< 3^{\text{rd}}$  percentile for gestational age.

**Results:** Low CPR was associated with increased risk of adverse perinatal outcome, presence of surrogate markers of perinatal hypoxia, cesarean section for presumed fetal distress in labor and birth of neonates with birthweight  $< 3^{\text{rd}}$  percentile. However, multivariable regression analysis demonstrated that the prediction of these adverse outcomes by maternal demographic characteristics and medical history was only marginally improved by the addition of CPR. The performance of low CPR in the prediction of each adverse outcome was poor, with detection rates of 13–26% and false positive rate of about 10%. In appropriate for gestational age (AGA) neonates with birthweight  $\geq 10^{\text{th}}$  percentile the predictive accuracy of CPR was low with positive and negative likelihood ratios (LRs) ranging from 1.21 to 1.82, and 0.92 to 0.98, respectively; although the accuracy was better in small for gestational age (SGA) neonates this was also low with positive LRs of 1.31 to 2.26 and negative LRs of 0.69 to 0.92. Similar values were obtained in fetuses classified as SGA and AGA according to the estimated fetal weight. In the prediction of adverse outcomes within two weeks, rather than at any stage, after assessment the detection rate was higher but this was achieved at higher false positive rate and therefore similar positive and negative LRs.

**Conclusion:** In pregnancies undergoing routine antenatal assessment at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation measurement of CPR provides poor prediction of adverse perinatal outcome in both SGA and AGA fetuses. Consequently, there is no justification in a shift of the focus of prenatal care from identification of pregnancies with low estimated fetal weight to that of pregnancies with low CPR.

## INTRODUCTION

In the 1980's studies of fetal blood obtained by cordocentesis from small for gestational age (SGA) fetuses demonstrated that increased impedance to flow, reflected in high pulsatility index (PI) in the umbilical artery (UA), and decreased PI in the fetal middle cerebral artery (MCA) are associated with fetal hypoxemia and acidemia.<sup>1-4</sup> It was subsequently shown that in SGA fetuses the cerebroplacental ratio (CPR) was a better predictor of adverse perinatal outcome than MCA-PI or UA-PI alone and that low CPR is associated with increased rates of perinatal death, cesarean section for fetal distress in labor, neonatal acidosis, 5 minute Apgar scores <7, and neonatal intensive care unit (NICU) stay >24 hours.<sup>5-8</sup> Renewed interest in the CPR has been stimulated by the possibility that this index may be predictive of adverse perinatal outcome not only in SGA but also in appropriately grown for gestational age (AGA) fetuses.<sup>9-12</sup> However, these studies have mainly examined high-risk pregnancies and did not report on the performance of CPR in the prediction of adverse outcome.

A screening study in 30,870 women with singleton pregnancies attending for a routine hospital visit at 30-34 weeks' gestation investigated the potential value of CPR in the prediction of adverse perinatal outcome and reported that although there was an association between CPR and birthweight Z-score, umbilical cord blood pH and admission to NICU, the performance of screening by CPR was poor with detection rates (DR) of 5-11% at false positive rate (FPR) of 5%.<sup>13</sup> A possible explanation for such poor performance of screening was that the perinatal adverse events at term were too remote from the gestational age at which CPR was assessed. However, another study of 6,178 singleton pregnancies routinely screened at 35-37 weeks' gestation, also reported significant associations between CPR and indicators of adverse perinatal outcome but again the performance of screening by CPR was poor with DR of 6-15%, at FPR of 6%.<sup>14</sup>

The objective of this extended study of 47,211 singleton pregnancies undergoing routine screening at 35<sup>+6</sup> - 36<sup>+6</sup> weeks' gestation is to investigate further the potential value of CPR in the prediction of adverse perinatal outcome.

## METHODS

### Study population

This was a prospective study in women with singleton pregnancies attending for a routine hospital visit at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation at King's College Hospital, London or Medway Maritime Hospital, Gillingham, UK between March 2014 and September 2018. This visit included recording of maternal demographic characteristics and medical history, ultrasound examination for fetal anatomy and measurement of fetal head circumference, abdominal circumference and femur length for calculation of EFW<sup>15,16</sup> and transabdominal color Doppler ultrasound for measurement of the UA-PI and MCA-PI.<sup>17</sup> Gestational age was determined by the measurement of fetal crown-rump length at 11-13 weeks.<sup>18</sup>

The women gave written informed consent to participate in the study, which was approved by the National Health Service Research Ethics Committee. The inclusion criteria for this study were singleton pregnancies examined at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation and delivering a non-malformed live birth or stillbirth. We excluded pregnancies with aneuploidies and major fetal abnormalities. Data from the first 6,178 pregnancies included in this study were reported previously.<sup>14</sup>

### Patient characteristics

Patient characteristics recorded included maternal age, racial origin (White, Black, South Asian, East Asian and mixed), method of conception (spontaneous or assisted by use of

1 ovulation induction drugs or *in vitro* fertilization), cigarette smoking during pregnancy,  
2 medical history of chronic hypertension or diabetes mellitus, obstetric history (nulliparous if  
3 no previous pregnancies at  $\geq 24$  weeks and parous with or without previous history of  
4 preeclampsia (PE) and / or birth of SGA neonate with birthweight  $< 10^{\text{th}}$  percentile) and  
5 presence of obstetric cholestasis or gestational diabetes mellitus in the current pregnancy.  
6 Maternal weight and height were measured and body mass index (BMI) was calculated.

### 7 Outcome measures

8  
9  
10 Data on pregnancy outcome were collected from the hospital maternity records. The  
11 following prespecified outcome measures were considered: first, adverse perinatal outcome  
12 consisting of stillbirth, neonatal death or hypoxic ischemic encephalopathy grades 2 and 3,  
13 second, presence of surrogate markers of perinatal hypoxia consisting of umbilical arterial or  
14 venous cord blood pH  $\leq 7$  and  $\leq 7.1$ , respectively, 5-minute Apgar score  $< 7$ , or admission to  
15 NICU for  $> 24$  hours, third, cesarean section for presumed fetal distress in labor, and fourth,  
16 SGA neonates with birthweight  $< 3^{\text{rd}}$  percentile.<sup>16</sup> Cesarean section for presumed fetal  
17 distress in labor was carried out if there was evidence of a pathological electronic fetal heart  
18 rate pattern, a STAN event on fetal electrocardiogram analysis or fetal scalp pH  $< 7.1$ .  
19 Hypoxic-ischemic encephalopathy was diagnosed when there was disturbed neurologic  
20 function with evidence of perinatal hypoxia reflected in either a 5-minute Apgar score  $< 5$  or  
21 umbilical artery cord pH  $< 7.0$  or base deficit  $> 12$  mmol/L, supported by neuroimaging  
22 evidence of acute brain injury.  
23  
24

### 25 Statistical analysis

26  
27  
28 Data were expressed as median (interquartile range [IQR]) for continuous variables and n  
29 (%) for categorical variables. Mann-Whitney U-test and  $\chi^2$ -square test or Fisher's exact test,  
30 were used for comparing outcome groups for continuous and categorical data, respectively.  
31 Significance was assumed at 5%.  
32

33  
34 Univariable and multivariable logistic regression analysis was carried out to determine which  
35 of the factors from maternal or pregnancy characteristics and measurements of UA-PI and  
36 MCA-PI and their ratio, provided a significant contribution in the prediction of each of the four  
37 outcome measures. Prior to the regression analysis, the continuous variables, such as age,  
38 weight and height were centred by subtracting the arithmetic mean from each value to avoid  
39 effects of multicollinearity. Multiple categorical variables were dummy coded as binary  
40 variables to estimate the independent effect of each category. The measured UA-PI and  
41 MCA-PI and their ratio were converted to multiples of the median (MoM) after adjustment for  
42 gestational age.<sup>17</sup> The birth weight Z-score was derived from the Fetal Medicine Foundation  
43 fetal and neonatal population weight charts.<sup>16</sup> We estimated cut-offs for the 90<sup>th</sup> percentile  
44 for UA-PI and 10<sup>th</sup> percentiles for MCA-PI and CPR and determined the prevalence of  
45 abnormal Doppler values in each of the outcome groups. The values of UA-PI  $> 90^{\text{th}}$   
46 percentile, MCA-PI  $< 10^{\text{th}}$  percentile and CPR  $< 10^{\text{th}}$  percentile were used as binary  
47 categorical variables in the multivariable regression analysis for each outcome measure.  
48 Predicted probabilities from logistic regression analysis were used to construct receiver  
49 operating characteristic (ROC) curves to assess performance of screening for these adverse  
50 outcomes. The area under ROC (AUROC) curves for fetal Doppler alone was compared to  
51 that obtained from all factors.<sup>17</sup> We examined the DR, FPR and positive and negative  
52 likelihood ratios (LR) of CPR  $< 10^{\text{th}}$  percentile for adverse perinatal outcome, presence of  
53 surrogate markers of perinatal hypoxia and cesarean section for presumed fetal distress in  
54 labor in the sub-groups of SGA (birthweight  $< 10^{\text{th}}$  percentile) and AGA (birthweight  $\geq 10^{\text{th}}$   
55 percentile) fetuses and neonates born within two weeks and at any stage after assessment.  
56  
57  
58

59 The statistical package SPSS 24.0 (IBM SPSS Statistics for Windows, Version 24.0,  
60 Armonk, NY: IBM Corp; 2016) was used for data analyses.  
61  
62  
63  
64  
65

## RESULTS

### Study population

During the study period, we prospectively examined and measured MCA-PI and UA-PI in 47,521 singleton pregnancies. We excluded 268 (0.6%) for major fetal abnormalities or genetic syndromes diagnosed prenatally or postnatally and 42 (0.1%) for no follow-up. The study population comprised 47,211 pregnancies. The median interval between assessment at 35<sup>+6</sup> – 36<sup>+6</sup> weeks' gestation and delivery was 3.7 (IQR 2.9, 4.7) weeks.

### Adverse perinatal outcome

Adverse perinatal outcome occurred in 130 (0.3%) cases and included 53 stillbirths, 11 neonatal deaths and 66 cases of HIE grades 2 or 3. The maternal and pregnancy characteristics of those with and without adverse perinatal outcome are compared in Table 1. In pregnancies with adverse perinatal outcome there was a higher median maternal weight and BMI, higher incidence of nulliparous women, lower incidence of parous women without previous SGA or PE, and lower median MoM values for MCA-PI and CPR.

Multivariable regression analysis demonstrated that in prediction of adverse perinatal outcome there was a statistically significant contribution from maternal BMI, nulliparity, MCA-PI and CPR <10<sup>th</sup> percentile ( $R^2=0.021$ ;  $p<0.001$ ; Table 2 and S1). The performance of screening by maternal factors alone in prediction of adverse perinatal outcome (DR 17.7% at FPR of 10%) was significantly improved by the addition of MCA and CPR (DR 26.2% at FPR of 10%; AUROC: 0.607, 95% CI 0.603, 0.612 vs. 0.644, 95% CI 0.639, 0.648;  $p=0.041$ ) (Figure 1).

### Surrogate markers of perinatal hypoxia

The 47,081 pregnancies without adverse perinatal outcome, included 1,370 (2.9%) with and 45,711 without surrogate markers of perinatal hypoxia. The maternal and pregnancy characteristics of these two groups are compared in Table S2. In pregnancies with surrogate markers of perinatal hypoxia there was a lower median maternal age, height, MCA-PI MoM, CPR MoM and birthweight, a lower incidence of women from East Asian and mixed racial origin, higher median maternal weight and BMI and higher incidence of cigarette smokers, women from Black racial origin, nulliparous women, those with diabetes mellitus, obstetric cholestasis, and birthweight <10<sup>th</sup> percentile.

Multivariable regression analysis demonstrated that in prediction of pregnancies with surrogate markers of perinatal hypoxia there was a statistically significant contribution from maternal BMI, cigarette smoking, Black and mixed racial origin, nulliparity, obstetric cholestasis, MCA-PI and CPR <10<sup>th</sup> percentile ( $R^2=0.021$ ;  $p<0.001$ ; Table 2 and S3). The performance of screening by maternal factors alone in prediction of adverse neonatal outcome (DR 17.2% at FPR of 10%) was significantly improved by the addition of MCA and CPR (DR 18.7% at FPR of 10%; AUROC: 0.588, 95% CI 0.583, 0.592 vs. 0.595, 95% CI 0.590, 0.599;  $p=0.032$ ) (Figure 1).

### Cesarean section for presumed fetal distress

The 47,158 pregnancies with livebirths, included 34,834 with vaginal delivery following spontaneous or induced labor, 5,475 with elective cesarean section for a variety of indications and 6,653 with cesarean section following spontaneous or induced labor; in the latter group, the indication for cesarean section was presumed fetal distress in 2,590 cases (Figure 2). Among those who underwent elective cesarean section ( $n=5,671$ ) there were a

1 variety of indications including breech or transverse lie, placenta previa, previous cesarean  
2 section or traumatic birth, maternal medical disorder or maternal request (n=5,475) and fetal  
3 compromise diagnosed by abnormal Doppler findings or fetal-heart rate patterns in SGA  
4 fetuses (n=196).

5 The maternal and pregnancy characteristics of those delivering by cesarean section for  
6 presumed fetal distress in labor are compared to those with vaginal delivery in Table S4. In  
7 pregnancies delivering by cesarean section for presumed fetal distress there was a lower  
8 median height, MCA-PI MoM, CPR MoM and birthweight, higher median maternal age,  
9 weight, BMI and UA-PI MoM and higher incidence of women of Black and South Asian racial  
10 origin, those who conceived by *in vitro* fertilisation, nulliparous women and parous women  
11 with a previous history of SGA or PE, chronic hypertension, diabetes mellitus, gestational  
12 diabetes, EFW <10<sup>th</sup> percentile and birthweight <10<sup>th</sup> percentile.

13  
14  
15 Multivariable regression analysis demonstrated that in prediction of cesarean section for  
16 presumed fetal distress in labor, there was a statistically significant contribution from  
17 maternal age, BMI, cigarette smoking, Black and South Asian racial origin, conception by *in*  
18 *vitro* fertilization nulliparity, previous PE or SGA, chronic hypertension, diabetes mellitus,  
19 EFW <10<sup>th</sup> percentile, and CPR <10<sup>th</sup> percentile (R<sup>2</sup>=0.087; p<0.0001; Table 2 and S5). In  
20 screening for cesarean section for presumed fetal distress by maternal factors alone the DR  
21 was 29.5% at FPR of 10%; addition of CPR did not improve the performance of screening  
22 (AUROC: 0.705, 95% CI 0.694, 0.705 vs. 0.706, 95% CI 0.695, 0.716; p=0.222) (Figure 1).

23  
24  
25 In SGA neonates delivered by elective cesarean section for presumed fetal distress, the  
26 performance of screening by maternal factors, obstetric and medical history (DR 85.2%,  
27 FPR 10%) was improved by the addition of UA-PI, MCA-PI and CPR (DR 91.8%, FPR 10%;  
28 AUROC: 0.896, 95% CI 0.868, 0.923 vs. 0.971, 95% CI 0.961, 0.981; p<0.0001). The CRP  
29 was <10<sup>th</sup> percentile in 67.9% (133/196) of cases with cesarean section for presumed fetal  
30 distress and in 9.5% (3307/34834) of those with vaginal delivery.

### 31 SGA neonates with birthweight <3<sup>rd</sup> percentile

32  
33  
34 The study population of 47,211 pregnancies included 2,102 (4.5%) with birthweight <3<sup>rd</sup>  
35 percentile and 45,109 (95.5%) with birthweight ≥3<sup>rd</sup> percentile. The maternal and pregnancy  
36 characteristics of those with and without SGA <3<sup>rd</sup> are compared in Table S6. In pregnancies  
37 with SGA <3<sup>rd</sup> percentile there was a lower median maternal age, weight, height, BMI, MCA-  
38 PI MoM, CPR MoM, and birthweight, lower incidence of women with diabetes mellitus,  
39 higher median UA-PI MoM and higher incidence of women of Black, South Asian and mixed  
40 racial origin, those who conceived by *in vitro* fertilisation, nulliparous women and those  
41 parous women with a previous history of SGA or PE, chronic hypertension, EFW <10<sup>th</sup>  
42 percentile and birthweight <10<sup>th</sup> percentile. The CRP was <10<sup>th</sup> percentile in 25.9% of cases  
43 with birthweight <3<sup>rd</sup> percentile and in 9.1% of those with birthweight ≥3<sup>rd</sup> percentile.

44  
45  
46 Multivariable regression analysis demonstrated that in prediction of SGA <3<sup>rd</sup> there was a  
47 statistically significant contribution from maternal age, BMI, Black, South Asian and mixed  
48 racial origin, cigarette smoking, diabetes mellitus, parity, UA-PI >90<sup>th</sup> percentile, MCA-PI  
49 <10<sup>th</sup> percentile and CPR <10<sup>th</sup> percentile (R<sup>2</sup>=0.335; p=0.001; Table 2 and S7). The  
50 performance of screening by maternal factors and EFW alone in prediction of SGA <3<sup>rd</sup> (DR  
51 68.7% at FPR of 10%) was improved by the addition of UA, MCA and CPR (DR 69.7% at  
52 FPR of 10%; AUROC: 858, 95% CI 849, 867 vs. 0.865, 95% CI 0.856, 0.874; p<0.0001)  
53 (Figure 1).

### 54 Performance of screening in pregnancies with SGA and AGA fetuses or neonates

55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 There was a significant association between  $\log_{10}$ MoM CPR and birthweight Z-score  
2 ( $r=0.210$ ,  $P<0.0001$ ). The incidence of CPR  $<10^{\text{th}}$  percentile increased with decreasing  
3 birthweight percentile; the incidence was 20.9% for birthweight  $<10^{\text{th}}$  percentile, 12.3%  
4 between the  $10^{\text{th}}$  and  $25^{\text{th}}$  percentiles, 9.8% between the  $25^{\text{th}}$  and  $50^{\text{th}}$  percentiles, 7.6%  
5 between the  $50^{\text{th}}$  and  $75^{\text{th}}$  percentiles, 6.0% between the  $75^{\text{th}}$  and  $90^{\text{th}}$  percentiles and 5.3%  
6 for birthweight  $>90^{\text{th}}$  percentile.

7  
8 The incidence of adverse perinatal outcome was 0.4% (20/5,509) in babies with birthweight  
9  $<10^{\text{th}}$  percentile and 0.3% (110/41,702) in those with birthweight  $\geq 10^{\text{th}}$  percentile ( $p=0.186$ ).  
10 Consequently, 84.6% (110/130) of adverse perinatal outcome occurred in AGA babies. The  
11 CRP was  $<10^{\text{th}}$  percentile in 20.0% of cases with and in 9.8% of those without adverse  
12 perinatal outcome. The incidence of surrogate markers of perinatal hypoxia was 4.2%  
13 (230/5,489) in babies with birthweight  $<10^{\text{th}}$  percentile and in 2.7% (1,140/41,592) of those  
14 with birthweight  $\geq 10^{\text{th}}$  percentile ( $p<0.0001$ ). Consequently, 83.3% (1,141/1,370) of adverse  
15 perinatal outcome occurred in AGA babies. The CRP was  $<10^{\text{th}}$  percentile in 13.7% of cases  
16 with and in 9.7% of those without surrogate markers of perinatal hypoxia. The incidence of  
17 cesarean section for presumed fetal distress in labor was 11.1% (503/4,543) in babies with  
18 birthweight  $<10^{\text{th}}$  percentile and in 6.3% (2,087/32,881) with birthweight  $\geq 10^{\text{th}}$  percentile  
19 ( $p<0.0001$ ). Consequently, 80.6% (2,087/2,590) of cesarean section for presumed fetal  
20 distress occurred in AGA babies. The CRP was  $<10^{\text{th}}$  percentile in 13.1% of cases with  
21 cesarean section for presumed fetal distress and in 9.5% of those with vaginal delivery.  
22  
23

24 The DR, FPR positive LR and negative LR of CPR  $<10^{\text{th}}$  percentile in the prediction of  
25 adverse perinatal outcome, perinatal hypoxia, cesarean section for presumed fetal distress  
26 in pregnancies with SGA and AGA fetuses and neonates are shown in Table 3. In AGA  
27 neonates the predictive accuracy of CPR was low with positive and negative LRs ranging  
28 from 1.21 to 1.82, and 0.92 to 0.98, respectively; although the accuracy was better in SGA  
29 neonates this was also low with positive LRs of 1.31 to 2.26 and negative LRs of 0.69 to  
30 0.92. Similar values were obtained in fetuses classified as SGA and AGA according to the  
31 estimated fetal weight. In the prediction of adverse outcomes within two weeks, rather than  
32 at any stage, after assessment the DR was higher but this was achieved at higher FPR and  
33 therefore similar positive and negative LRs.  
34  
35

## 36 COMMENT

### 37 Principal findings of the study

38  
39 The findings of this study of routine ultrasound examination in singleton pregnancies at  $35^{+0}$ -  
40  $36^{+6}$  weeks' gestation demonstrate that although the incidence of adverse perinatal  
41 outcome, presence surrogate markers of perinatal hypoxia and cesarean section for  
42 presumed fetal distress in labor is higher in pregnancies with SGA compared to AGA  
43 fetuses; 80-85% of these adverse events occur in the AGA group. If it was to be assumed  
44 that first, these adverse outcomes are the consequence of impaired placentation and fetal  
45 hypoxia and second, low CPR is a good marker of fetal hypoxia irrespective of fetal size, it  
46 should be anticipated that low CPR would be a good predictor of adverse outcome. It could  
47 then be argued that prenatal care should be directed at identifying hypoxemic rather than  
48 small fetuses and, consequently, screening should focus on the detection of pregnancies  
49 with low CPR rather than those with low estimated fetal weight.  
50  
51  
52  
53

54 We found that low CPR was associated with increased risk of adverse perinatal outcome,  
55 presence of surrogate markers of perinatal hypoxia, cesarean section for presumed fetal  
56 distress in labor and birth of neonates with birthweight  $<3^{\text{rd}}$  percentile. However,  
57 multivariable regression analysis demonstrated that the prediction of these adverse  
58 outcomes by maternal demographic characteristics and medical history was only marginally  
59 improved by the addition of CPR. The performance of low CPR in the prediction of each  
60  
61  
62  
63  
64  
65



1 adverse outcome was poor, with DR of 13–26% and FPR of about 10%. The DR of adverse  
2 outcomes was higher in SGA than in AGA babies, irrespective of whether the classification  
3 was based on estimated fetal weight or birthweight and in pregnancies delivering within two  
4 weeks rather than at any stage after assessment. However, such increase in DR was  
5 accompanied by an increase in FPR and the predictive accuracy of the test was low,  
6 reflected in low positive LRs and high negative LRs irrespective of fetal size or interval  
7 between testing and delivery.

8  
9 The low performance of CPR in the prediction of adverse perinatal outcomes suggests that  
10 either CPR provides poor assessment of fetal oxygenation or that first, most cases of  
11 stillbirth at term are not associated with impaired placentation and chronic fetal hypoxia and  
12 second, the contribution of maternal and pregnancy characteristics as well as events in labor  
13 play a much greater role than prelabor fetal oxygenation in the development of fetal distress  
14 in labor or adverse neonatal outcome.

### 15 16 Strengths and limitations of the study

17  
18 The strengths of our study are first, examination of a large number of pregnancies, including  
19 5,509 that delivered SGA neonates, attending for routine assessment of fetal growth and  
20 wellbeing at a prespecified gestational-age range at the end of the third trimester of  
21 pregnancy, second, measurement of MCA-PI and UA-PI by appropriately-trained doctors,  
22 and third, use of a wide range of well accepted indicators for adverse perinatal outcome.

23  
24 The main limitation of this and most previous studies investigating the value of CPR in the  
25 prediction of adverse pregnancy outcome is that the results of the ultrasound scan were  
26 made available to the attending obstetricians who would have taken specific actions of  
27 further monitoring and planned delivery of the cases with suspected SGA and fetal  
28 compromise. In our study 196 such pregnancies had elective delivery by cesarean section;  
29 had this not been carried out it is possible that some of the cases would have resulted in  
30 stillbirth, cesarean section for fetal distress in labor and birth asphyxia. Consequently, the  
31 performance of screening by CPR for adverse perinatal outcome in SGA fetuses would have  
32 been negatively biased.

### 33 34 Comparison with findings from previous studies

35  
36 Several prospective and retrospective studies in small numbers of third trimester high-risk  
37 pregnancies reported an association between low CPR or low MCA-PI and increased risk of  
38 adverse perinatal outcomes, but they did not report on the predictive performance of the  
39 test.<sup>7</sup>

40  
41 Our study, in 47,211 pregnancies, evaluated CPR at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation as part of  
42 routine screening for adverse perinatal outcome in all pregnant women, irrespective of fetal  
43 size or interval from delivery. Our findings confirm the association between low CPR and  
44 adverse perinatal outcomes but demonstrate that the predictive performance of the test in  
45 both SGA and AGA fetuses is poor. These findings are consistent with those of our previous  
46 study in 30,870 pregnancies undergoing routine screening at 30-34 weeks' gestation.<sup>13</sup>

### 47 48 Implications for clinical practice

49  
50 About 85% of SGA neonates are born at term,<sup>20</sup> and in such neonates the risk of adverse  
51 outcome is substantially higher than in AGA neonates.<sup>21,22</sup> The traditional approach of  
52 identifying pregnancies with SGA fetuses is maternal abdominal palpation and serial  
53 measurements of symphysial-fundal height, which is advocated by national guidelines in the  
54 USA and many other developed countries; however, the predictive performance of such  
55 screening is poor.<sup>23-25</sup> There is some evidence that substantially improved prediction of SGA  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 is achieved by universal sonographic fetal biometry during the third trimester, especially at  
2 about 36 weeks' gestation.<sup>20,26-30</sup> A prospective study in 19,208 singleton pregnancies  
3 undergoing routine antenatal assessment at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation reported that a  
4 combination of maternal demographic characteristics and medical history with  
5 sonographically estimated fetal weight predicted 90% of SGA neonates delivering within two  
6 weeks and at any stage after assessment at respective screen positive rates of about 20%  
7 and 30%.<sup>20</sup>

8  
9 In this study we found that in pregnancies undergoing such routine antenatal assessment at  
10 35<sup>+0</sup> - 36<sup>+6</sup> weeks measurement of CPR provides poor prediction of adverse perinatal  
11 outcome in both SGA and AGA fetuses. Consequently, there is no justification for a shift of  
12 the focus of prenatal care from identification of pregnancies with low estimated fetal weight  
13 to that of pregnancies with low CPR. There is no evidence that incorporating CPR in the  
14 management of AGA fetuses reduces perinatal death or other adverse perinatal outcomes,  
15 but there is a risk that such practice would increase early iatrogenic delivery.  
16

17  
18 There is also lack of evidence that incorporating CPR in the management of SGA fetuses  
19 reduces perinatal death or other adverse perinatal outcomes. This raises the question as to  
20 the best management of pregnancies at high risk of delivering SGA neonates; specifically,  
21 30% of the population identified by combined screening with maternal factors and fetal  
22 biometry at 35<sup>+0</sup> - 36<sup>+6</sup> weeks' gestation to contain 90% of SGA neonates.<sup>20</sup> On the basis of  
23 results from observational studies measurement of CPR can contribute in the differentiation  
24 of constitutionally-small from growth-restricted fetuses.<sup>8,30</sup> It could then be argued that in the  
25 subgroup of the 30% of the population at high risk of delivering SGA neonates those with  
26 CPR <10<sup>th</sup> percentile should undergo iatrogenic delivery at around 37 weeks, whereas those  
27 with CPR ≥10<sup>th</sup> percentile could have close surveillance with delayed delivery until 39-40  
28 weeks. The extent to which such policy would reduce adverse perinatal outcome merits  
29 further investigation.  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

1. Nicolaides KH, Soothill PW, Rodeck CH, Campbell S. Ultrasound guided sampling of umbilical cord and placental blood to assess fetal wellbeing. *Lancet* 1986;1:1065-7.
2. Soothill PW, Nicolaides KH, Campbell S. Prenatal asphyxia, hyperlacticaemia, hypoglycaemia and erythroblastosis in growth retarded fetuses. *BMJ* 1987;294:1051-3.
3. Nicolaides KH, Bilardo KM, Soothill PW, Campbell S. Absence of end diastolic frequencies in the umbilical artery a sign of fetal hypoxia and acidosis. *BMJ* 1988;297:1026-7.
4. Vyas S, Nicolaides KH, Bower S, Campbell S. Middle cerebral artery flow velocity waveforms in fetal hypoxaemia. *Br J Obstet Gynaecol* 1990;97:797-803.
5. Bahado-Singh RO, Kovanci E, Jeffres A, et al. The Doppler cerebroplacental ratio and perinatal outcome in intrauterine growth restriction. *Am J Obstet Gynecol* 1999;180:750-6.
6. Gramellini D, Folli MC, Raboni S, Vadora E, Merialdi A. Cerebral-umbilical Doppler ratio as a predictor of adverse perinatal outcome. *Obstet Gynecol* 1992;79:416-20.
7. DeVore GR. The importance of the cerebroplacental ratio in the evaluation of fetal well-being in SGA and AGA fetuses. *Am J Obstet Gynecol* 2015;213: 5-15.
8. Flood K, Unterscheider J, Daly S, et al. The role of brain sparing in the prediction of adverse outcomes in intrauterine growth restriction: results of the multicenter PORTO Study. *Am J Obstet Gynecol* 2014;211:288.e1-5.
9. Prior T, Mullins E, Bennett P, Kumar S. Prediction of intrapartum fetal compromise using the cerebroumbilical ratio: a prospective observational study. *Am J Obstet Gynecol* 2013;208:124.e1-6.
10. Khalil AA, Morales-Rosello J, Morlando M, et al. Is fetal cerebroplacental ratio an independent predictor of intrapartum fetal compromise and neonatal unit admission? *Am J Obstet Gynecol* 2015;213:54.e1-10.
11. Khalil AA, Morales-Rosello J, Elsadig M, et al. The association between fetal Doppler and admission to neonatal unit at term. *Am J Obstet Gynecol* 2015;213:57.e1-7.
12. Khalil A, Morales-Rosello J, Khan N, et al. Is cerebroplacental ratio a marker of impaired fetal growth velocity and adverse pregnancy outcome? *Am J Obstet Gynecol* 2017;216:606.e1-606.e10.
13. Bakalis S, Akolekar R, Gallo DM, Poon LC, Nicolaides KH. Umbilical and fetal middle cerebral artery Doppler at 30-34 weeks' gestation in the prediction of adverse perinatal outcome. *Ultrasound Obstet Gynecol* 2015;45:409-20.
14. Akolekar R, Syngelaki A, Gallo DM, Poon LC, Nicolaides KH. Umbilical and fetal middle cerebral artery Doppler at 35-37 weeks' gestation in the prediction of adverse perinatal outcome. *Ultrasound Obstet Gynecol* 2015;46:82-92.
15. Hammami A, Mazer Zumaeta A, Syngelaki A, Akolekar R, Nicolaides KH. Ultrasonographic estimation of fetal weight: development of new model and assessment of performance of previous models. *Ultrasound Obstet Gynecol*

2018;52:35-43.

16. Nicolaides KH, Wright D, Syngelaki A, Wright A, Akolekar R. Fetal Medicine Foundation fetal and neonatal population weight charts. *Ultrasound Obstet Gynecol* 2018;52:44-51.
17. Ciobanu A, Wright A, Syngelaki A, Wright D, Akolekar R, Nicolaides KH. Fetal Medicine Foundation reference ranges for umbilical artery and middle cerebral artery pulsatility index and cerebroplacental ratio. *Ultrasound Obstet Gynecol* 2018;doi: 10.1002/uog.20157.
18. Robinson HP, Fleming JE. A critical evaluation of sonar crown rump length measurements. *Br J Obstet Gynaecol* 1975;82:702-10.
19. Zweig MH, Campbell G. Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993; 39: 561-577.
20. Ciobanu A, Rouvali A, Syngelaki A, Akolekar R, Nicolaides KH. Prediction of small for gestational age neonates: Screening by maternal factors and biomarkers at 35-37 weeks' gestation. *Am J Obstet Gynecol*.
21. Mendez-Figueroa H, Truong VT, Pedroza C, Khan AM, Chauhan SP. Small-for-gestational-age infants among uncomplicated pregnancies at term: a secondary analysis of 9 Maternal-Fetal Medicine Units Network studies. *Am J Obstet Gynecol* 2016;215:628.e1-628.e7.
22. Madden JV, Flatley CJ, Kumar S. Term small-for-gestational-age infants from low-risk women are at significantly greater risk of adverse neonatal outcomes. *Am J Obstet Gynecol* 2018;218:525.e1-525.e9.
23. Bais JMJ, Eskes M, Pel M, Bonse GJ, Bleker OP. Effectiveness of detection of intrauterine retardation by abdominal palpation as screening test in a low-risk population: an observational study. *Eur J Obstet Gynecol Reprod Biol* 2004;116:164-9.
24. Lindhard A, Nielsen PV, Mouritsen LA, Zachariassen A, Sørensen HU, Rosenø H. The implications of introducing the symphyseal-fundal height-measurement. A prospective randomized controlled trial. *Br J Obstet Gynaecol* 1990;97:675-80.
25. McCowan LM, Figueras F, Anderson NH. Evidence-based national guidelines for the management of suspected fetal growth restriction: comparison, consensus, and controversy. *Am J Obstet Gynecol* 2018;218:S855-S868.
26. David C, Tagliavini G, Pilu G, Rudenholz A, Bovicelli L. Receiver-operator characteristic curves for the ultrasonographic prediction of small-for-gestational-age fetuses in low-risk pregnancies. *Am J Obstet Gynecol* 1996;174:1037-42.
27. Bakalis S, Silva M, Akolekar R, Poon LC, Nicolaides KH. Prediction of small for gestational age neonates: screening by fetal biometry at 30–34 weeks. *Ultrasound Obstet Gynecol* 2015;45:551-8.
28. Sovio U, White IR, Dacey A, Pasupathy D, Smith GCS. Screening for fetal growth restriction with universal third trimester ultrasonography in nulliparous women in the Pregnancy Outcome Prediction (POP) study: a prospective cohort study. *Lancet* 2015;386:2089-97.

- 1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65
29. Gaccioli F, Aye ILMH, Sovio U, Charnock-Jones DS, Smith GCS. Screening for fetal growth restriction using fetal biometry combined with maternal biomarkers. *Am J Obstet Gynecol* 2018;218:S725-37.
30. Figueras F, Caradeux J, Crispi F, Eixarch E, Peguero A, Gratacos E. Diagnosis and surveillance of late-onset fetal growth restriction. *Am J Obstet Gynecol* 2018;218:S790-S802.e1.

**Figure legend**

**Figure 1.** Receiver operating characteristic plots of screening for adverse perinatal outcome (a), surrogate markers of perinatal hypoxia (b), cesarean section for presumed fetal distress in labor (c), elective cesarean section for presumed fetal distress in small for gestational age fetuses (d) and birthweight <3<sup>rd</sup> percentile (e) by maternal factors (black curve) and the combination of maternal factors and Doppler findings (red curve).

**Figure 2.** Flow chart on the indications and method of delivery of the study population.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1.** Maternal and pregnancy characteristics in pregnancies with and without adverse perinatal outcome.

Maternal and pregnancy characteristics	No adverse outcome (n=47,081)	Adverse outcome (n=130)
Maternal age in years, median (IQR)	31.6 (27.3-35.4)	31.1 (27.3-34.9)
Maternal weight in kg, median (IQR)	79.0 (70.8-90.0)	83.0 (73.4-92.0)*
Maternal height in cm, median (IQR)	165 (160-169)	165 (161-169)
Maternal body mass index in kg/m <sup>2</sup> , median (IQR)	29.1 (26.2-32.9)	29.7 (27.3-34.5)*
Cigarette smoker, n (%)	3,840 (8.2)	11 (8.5)
Racial origin		
White, n (%)	34,994 (74.3)	92 (70.8)
Black, n (%)	7,461 (15.8)	28 (21.5)
South Asian, n (%)	2,250 (4.8)	4 (3.1)
East Asian, n (%)	966 (2.1)	2 (1.5)
Mixed, n (%)	1,410 (3.0)	4 (3.1)
Conception		
Natural, n (%)	45,465 (96.6)	127 (97.7)
Use of ovulation induction drugs, n (%)	264 (0.6)	0
<i>In vitro</i> fertilization, n (%)	1,352 (2.9)	3 (2.3)
Obstetric history		
Nulliparous, n (%)	21,389 (45.4)	76 (58.5)**
Parous, previous SGA or PE, n (%)	4,216 (9.0)	6 (4.6)
Parous, no previous SGA or PE, n (%)	21,476 (45.6)	48 (36.9)*
Medical disorders		
Chronic hypertension, n (%)	595 (1.3)	1 (0.8)
Diabetes mellitus, n (%)	381 (0.8)	0
Pregnancy complications		
Gestational diabetes, n (%)	2029 (4.3)	4 (3.1)
Obstetric cholestasis, n (%)	496 (1.1)	2 (1.5)
Doppler indices		
Umbilical artery PI in MoM, median (IQR)	1.01 (0.91-1.11)	1.01 (0.91-1.10)
Umbilical artery PI >90 <sup>th</sup> percentile, n (%)	4,090 (8.7)	10 (7.7)
Middle cerebral artery PI in MoM, median (IQR)	1.00 (0.90-1.10)	0.95 (0.85-1.06)**
Middle cerebral artery PI <10 <sup>th</sup> percentile, n (%)	3,984 (8.5)	25 (19.2)**
Cerebroplacental ratio in MoM, median (IQR)	0.99 (0.87-1.13)	0.96 (0.80-1.13)*
Cerebroplacental ratio <10 <sup>th</sup> percentile, n (%)	4,614 (9.8)	26 (20.0)**
Stillbirth (n=53)	-	13 (24.5)
Neonatal death (n=11)	-	2 (18.2)
Hypoxic ischemic encephalopathy (n=66)	-	11 (16.7)
Estimated weight <10 <sup>th</sup> percentile, n (%)	4,276 (9.1)	6 (4.6)
GA at delivery in weeks, median (IQR)	40.0 (39.0-40.9)	39.8 (38.7-41.0)
Birth weight in g, median (IQR)	3,420 (3100-3470)	3,360 (3075-3765)
Birth weight <10 <sup>th</sup> percentile, n (%)	5,489 (11.7)	20 (15.4)

IQR = interquartile range; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median

Significance value \* p<0.05; \*\* p<0.01

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 2.** Multivariable logistic regression analysis in prediction of adverse perinatal outcome, surrogate markers of perinatal hypoxia, cesarean section for fetal distress in labor and birthweight <3<sup>rd</sup> percentile from maternal and pregnancy characteristics.

Maternal and pregnancy characteristics	Adverse perinatal outcome		Perinatal hypoxia		CS for fetal distress		Birthweight <3 <sup>rd</sup> percentile	
	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value	OR (95% CI)	P value
Maternal age – 30 (years)					1.04 (1.03-1.04)	<0.0001	1.02 (1.01-1.03)	<0.0001
Maternal BMI – 30 (kg/m <sup>2</sup> )	1.05 (1.02-1.08)	<0.0001	1.04 (1.03-1.05)	<0.0001	1.07 (1.06-1.08)	<0.0001	0.98 (0.97-0.99)	0.001
Cigarette smoker			1.41 (1.19-1.68)	<0.0001	1.40 (1.20-1.63)	<0.0001	2.38 (2.05-2.76)	<0.0001
Racial origin								
Black			1.17 (1.02-1.35)	0.029	1.90 (1.71-2.10)	<0.0001	2.04 (1.79-2.32)	<0.0001
South Asian					1.57 (1.32-1.87)	<0.0001	2.27 (1.90-2.71)	<0.0001
Mixed			0.61 (0.41-0.90)	0.014			1.72 (1.32-2.24)	<0.0001
Conception								
<i>In vitro</i> fertilization					1.34 (1.08-1.67)	0.009		
Obstetric history								
Nulliparous	1.71 (1.20-2.43)	0.003	1.42 (1.28-1.58)	<0.0001	3.92 (3.54-4.33)	<0.0001	2.34 (2.08-2.64)	<0.0001
Parous, previous PE or SGA					1.51 (1.27-1.80)	<0.0001	2.54 (2.18-2.97)	<0.0001
Medical complications								
Chronic hypertension					1.52 (1.13-2.06)	0.007		
Diabetes mellitus					1.70 (1.14-2.54)	0.010	0.44 (0.21-0.94)	0.033
Pregnancy complications								
Gestational diabetes								
Cholestasis			1.68 (1.12-2.53)	0.012				
Estimated fetal weight <10 <sup>th</sup> percentile					1.25 (1.09-1.43)	0.001	20.02 (18.07-22.19)	<0.0001
Doppler indices								
UA-PI >90 <sup>th</sup> percentile							1.67 (1.44-1.94)	<0.0001
MCA-PI <10 <sup>th</sup> percentile	1.97 (1.20-3.23)	0.007	1.26 (1.05-1.50)	0.014			1.58 (1.35-1.85)	<0.0001
CPR <10 <sup>th</sup> percentile	1.71 (1.05-2.79)	0.031	1.36 (1.15-1.61)	<0.0001	1.31 (1.16-1.48)	<0.0001	1.67 (1.43-1.95)	<0.0001

OR = odds ratio; CI = confidence interval; BMI = body mass index; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

**Table 3.** Predictive performance of cerebroplacental ratio <10<sup>th</sup> percentile for adverse perinatal outcome, surrogate markers of perinatal hypoxia and cesarean section for fetal distress in labor in small and appropriate for gestational age fetuses and neonates

Classification according to estimated fetal weight	Cerebroplacental ratio <10 <sup>th</sup> percentile			
	At any stage after assessment		Within two weeks of assessment	
	Weight ≥10 <sup>th</sup> percentile	Weight <10 <sup>th</sup> percentile	Weight ≥10 <sup>th</sup> percentile	Weight <10 <sup>th</sup> percentile
<b>Adverse perinatal outcome (n=130)</b>				
Detection rate	23/124 (18.5; 11.8, 25.2)	3/6 (50.0; 41.4, 58.6)	6/23 (26.1; 8.9, 43.3)	1/1
False positive rate	3,790/42,805 (8.9; 8.6, 9.2)	824/4,276 (19.3; 18.9, 19.7)	612/4,896 (12.5; 11.6, 13.4)	437/1,296 (33.7; 32.4, 35.0)
Positive likelihood ratio	2.08 (1.94, 2.22)	2.59 (2.44, 2.74)	2.09 (1.69, 2.49)	2.97 (2.49, 3.45)
Negative likelihood ratio	0.89 (0.80, 0.98)	0.62 (0.54, 0.70)	0.84 (0.58, 1.10)	0.00
<b>Perinatal hypoxia (n=1,370)</b>				
Detection rate	131/1,223 (10.7; 9.1, 12.3)	57/147 (38.8; 36.2, 41.4)	49/258 (19.0; 14.2, 23.8)	42/73 (57.5; 46.3, 68.6)
False positive rate	3,659/41,582 (8.8; 8.5, 9.1)	767/4,129 (18.6; 18.2, 19.0)	563/4,638 (12.0; 11.0, 12.9)	395/1,223 (32.3; 29.7, 34.9)
Positive likelihood ratio	1.22 (1.12, 1.32)	2.09 (1.95, 2.23)	1.58 (1.22, 1.94)	1.78 (1.04, 2.52)
Negative likelihood ratio	0.98 (0.89, 1.00)	0.75 (0.67, 0.83)	0.92 (0.65, 1.19)	0.63 (0.19, 1.07)
<b>Cesarean section for fetal distress (n=2,590)</b>				
Detection rate	261/2,296 (11.4; 10.2, 12.6)	79/294 (26.9; 25.2, 28.6)	51/213 (23.9; 18.2, 29.6)	55/121 (45.5; 36.6, 54.4)
False positive rate	2,751/31,576 (8.7; 8.4, 9.0)	556/3,258 (17.1; 16.7, 17.5)	393/3,320 (11.8; 10.7, 12.9)	253/861 (29.4; 26.4, 32.4)
Positive likelihood ratio	1.31 (1.20, 1.42)	1.57 (1.45, 1.69)	2.03 (1.55, 2.51)	1.55 (0.72, 2.38)
Negative likelihood ratio	0.97 (0.88, 1.00)	0.88 (0.79, 0.97)	0.86 (0.55, 1.17)	0.77 (0.19, 1.35)
<b>Classification according to birthweight</b>				
<b>Adverse perinatal outcome (n=130)</b>				
Detection rate	17/110 (15.5; 9.3, 21.74)	9/20 (45.0; 36.5, 53.6)	4/19 (21.0; 14.0, 28.1)	3/5 (60.0; 51.6, 68.4)
False positive rate	3,523/41,592 (8.5; 8.2, 8.8)	1,091/5,489 (19.9; 19.5, 20.3)	575/4,781 (12.0; 11.7, 12.3)	474/1,411 (33.6; 32.2, 35.1)
Positive likelihood ratio	1.82 (1.69, 1.95)	2.26 (2.51, 2.81)	1.75 (1.38, 2.12)	1.79 (1.10, 2.48)
Negative likelihood ratio	0.92 (0.83, 1.00)	0.69 (0.61, 0.77)	0.90 (0.63, 1.17)	0.60 (0.20, 1.00)
<b>Perinatal hypoxia (n=1,370)</b>				
Detection rate	116/1140 (10.2; 8.6, 11.8)	72/230 (31.3; 28.8, 33.8)	42/241 (17.4; 15.1, 19.8)	49/90 (54.4; 48.8, 60.0)
False positive rate	3,407/40,452 (8.4; 8.1, 8.7)	1,019/5,259 (19.4; 19.0, 19.8)	533/4,540 (11.7; 11.4, 12.0)	425/1,321 (32.2; 30.8, 33.7)
Positive likelihood ratio	1.21 (1.11, 1.31)	1.61 (1.49, 1.73)	1.49 (1.14, 1.84)	1.69 (1.00, 2.39)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49

6 Negative likelihood ratio	0.98 (0.89, 1.00)	0.85 (0.76, 0.94)	0.94 (0.66, 1.22)	0.67 (0.23, 1.11)
7 Cesarean section for fetal distress (n=2,590)				
8 Detection rate	220/2,087 (10.5; 9.3, 11.7)	120/503 (23.9; 22.3, 25.5)	45/206 (21.8; 20.0, 23.6)	61/128 (47.7; 43.3, 52.1)
9 False positive rate	2,573/30,794 (8.4; 8.0, 8.7)	734/4,040 (18.2; 17.8, 18.6)	368/3,250 (11.3; 11.0, 11.6)	278/931 (29.9; 28.5, 31.3)
10 Positive likelihood ratio	1.25 (1.14, 1.36)	1.31 (1.20, 1.42)	1.93 (1.36, 2.50)	1.60 (0.92, 2.28)
11 Negative likelihood ratio	0.98 (0.89, 1.00)	0.92 (0.83, 1.00)	0.88 (0.49, 1.27)	0.75 (0.28, 1.22)

**Table S1.** Univariable and multivariable logistic regression analysis in prediction of adverse perinatal outcome from maternal and pregnancy characteristics.

Maternal and pregnancy characteristics	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Maternal age – 30 (years)	0.99 (0.96-1.02)	0.536		
Maternal BMI – 30 (kg/m <sup>2</sup> )	1.05 (1.02-1.07)	0.002	1.05 (1.02-1.08)	<0.0001
Cigarette smoker	1.04 (0.56-1.93)	0.899		
Racial origin				
White	1.00 (Reference)			
Black	1.43 (0.93-2.18)	0.100		
South Asian	0.68 (0.25-1.84)	0.44		
East Asian	0.79 (0.19-3.20)	0.738		
Mixed	1.08 (0.40-3.94)	0.882		
Conception				
Natural	1.00 (Reference)			
Use of ovulation induction drugs	-	-		
<i>In vitro</i> fertilization	0.79 (0.25-2.50)	0.694		
Obstetric history				
Parous, no previous PE or SGA	1.00 (Reference)			
Nulliparous	1.59 (1.11-2.28)	0.012	1.71 (1.20-2.43)	0.003
Parous, previous PE or SGA	0.64 (0.27-1.49)	0.298		
Pregnancy complications				
Gestational diabetes	0.71 (0.26-1.91)	0.492		
Cholestasis	1.47 (0.36-5.95)	0.591		
Estimated fetal weight				
Z-score	1.07 (0.91-1.26)	0.427		
<10 <sup>th</sup> percentile	0.48 (0.21-1.10)	0.083		
Doppler indices				
UA-PI >90 <sup>th</sup> percentile	0.88 (0.46-1.67)	0.688		
MCA-PI <10 <sup>th</sup> percentile	2.58 (1.66-3.99)	<0.001	1.97 (1.20-3.23)	0.007
CPR <10 <sup>th</sup> percentile	2.30 (1.50-3.54)	<0.001	1.71 (1.05-2.79)	0.031

BMI = body mass index; OR = odds ratio; CI = confidence interval; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median.

**Table S2.** Maternal and pregnancy characteristics in pregnancies with and without surrogate markers of perinatal hypoxia.

Maternal and pregnancy characteristics	No surrogate markers (n=45,711)	Surrogate markers (n=1,370)
Maternal age in years, median (IQR)	31.6 (27.0-35.4)	30.9 (26.5-35.1)**
Maternal weight in kg, median (IQR)	79.0 (70.7-90.0)	81.0 (72.0-93.3)**
Maternal height in cm, median (IQR)	165 (160-169)	164 (160-168)**
Maternal body mass index in kg/m <sup>2</sup> , median (IQR)	29.1 (26.2-32.9)	30.1 (26.9-34.7)**
Cigarette smoker, n (%)	3692 (8.1)	148 (10.8)**
Racial origin		
White, n (%)	33974 (74.3)	1020 (74.5)
Black, n (%)	7213 (15.8)	248 (18.1)*
South Asian, n (%)	2191 (4.8)	59 (4.3)
East Asian, n (%)	949 (2.1)	17 (1.2)*
Mixed, n (%)	1384 (3.0)	26 (1.9)*
Conception		
Natural, n (%)	44147 (96.6)	1318 (96.2)
Use of ovulation induction drugs, n (%)	256 (0.6)	8 (0.6)
<i>In vitro</i> fertilization, n (%)	1308 (2.9)	44 (3.2)
Obstetric history		
Nulliparous, n (%)	20680 (45.2)	709 (51.8)**
Parous, previous SGA or PE, n (%)	4087 (8.9)	125 (9.1)
Parous, no previous SGA or PE, n (%)	20944 (45.8)	536 (39.1)
Medical disorders		
Chronic hypertension, n (%)	570 (1.2)	25 (1.8)
Diabetes mellitus, n (%)	360 (0.8)	22 (1.6)**
Pregnancy complications		
Gestational diabetes, n (%)	1960 (4.3)	68 (5.0)
Obstetric cholestasis, n (%)	471 (1.0)	25 (1.8)**
Doppler indices		
Umbilical artery PI in MoM, median (IQR)	1.01 (0.91-1.11)	1.02 (0.91-1.12)
Umbilical artery PI >90 <sup>th</sup> percentile, n (%)	3951 (8.6)	141 (10.3)
Middle cerebral artery PI in MoM, median (IQR)	1.00 (0.90-1.10)	0.99 (0.89-1.09)**
Middle cerebral artery PI <10 <sup>th</sup> percentile, n (%)	3831 (8.4)	153 (11.2)**
Cerebroplacental ratio in MoM, median (IQR)	0.99 (0.87-1.13)	0.99 (0.87-1.13)**
Cerebroplacental ratio <10 <sup>th</sup> percentile, n (%)	4426 (9.7)	188 (13.7)**
Estimated weight <10 <sup>th</sup> percentile, n (%)	4129 (9.0)	147 (10.7)
GA at delivery in weeks, median (IQR)	40.0 (39.0-40.9)	39.8 (38.3-40.9)**
Birth weight in g, median (IQR)	3420 (3105-3740)	3380 (3000-3755)**
Birth weight <10 <sup>th</sup> percentile, n (%)	5259 (11.5)	229 (16.7)**

IQR = interquartile range; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median

Significance value \* p<0.05; \*\* p<0.01

**Table S3.** Univariable and multivariable logistic regression analysis in prediction of surrogate markers of perinatal hypoxia from maternal and pregnancy characteristics.

Maternal and pregnancy characteristics	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Maternal age – 30 (years)	0.99 (0.98-1.00)	0.004		
Maternal BMI – 30 (kg/m <sup>2</sup> )	1.04 (1.03-1.05)	<0.0001	1.04 (1.03-1.05)	<0.0001
Cigarette smoker	1.36 (1.14-1.61)	<0.0001	1.41 (1.19-1.68)	<0.0001
Racial origin				
White	1.00 (Reference)			
Black	1.14 (0.99-1.31)	0.067	1.17 (1.02-1.35)	0.029
South Asian	0.88 (0.68-1.15)	0.348		
East Asian	0.60 (0.38-0.96)	0.034		
Mixed	0.60(0.40-0.88)	0.010	0.61 (0.41-0.90)	0.014
Conception				
Natural	1.00 (Reference)			
Use of ovulation induction drugs	1.00 (0.49-2.02)	0.995		
<i>In vitro</i> fertilization	1.10 (0.81-1.49)	0.543		
Obstetric history				
Parous, no previous PE or SGA	1.00 (Reference)			
Nulliparous	1.39 (1.24-1.55)	<0.0001	1.42 (1.28-1.58)	<0.0001
Parous, previous PE or SGA	1.20 (0.99-1.46)	0.069		
Medical complications				
Chronic hypertension	1.40 (0.94-2.10)	0.100		
Diabetes mellitus	1.87 (1.20-2.91)	0.006		
Pregnancy complications				
Gestational diabetes	1.14 (0.90-1.46)	0.281		
Cholestasis	1.70 (1.13-2.55)	0.010	1.68 (1.12-2.53)	0.012
Estimated fetal weight				
Z-score	1.02 (0.97-1.07)	0.566		
<10 <sup>th</sup> percentile	1.16 (0.97-1.38)	0.098		
Doppler indices				
UA-PI >90 <sup>th</sup> percentile	1.16 (0.97-1.38)	0.099		
MCA-PI <10 <sup>th</sup> percentile	1.45 (1.23-1.71)	<0.0001	1.26 (1.05-1.50)	0.014
CPR <10 <sup>th</sup> percentile	1.50 (1.29-1.75)	<0.0001	1.36 (1.15-1.61)	<0.0001

BMI = body mass index; OR = odds ratio; CI = confidence interval; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median.

**Table S4.** Maternal and pregnancy characteristics in pregnancies delivering by cesarean section for fetal distress in labor compared to those that delivered vaginally.

Maternal and pregnancy characteristics	Vaginal delivery (n=34,834)	Cesarean section (n=2,590)
Maternal age in years, median (IQR)	31.1 (26.8-34.9)	31.4 (27.1-35.3)*
Maternal weight in kg, median (IQR)	78.3 (70.0-89.0)	81.0 (72.0-92.2)**
Maternal height in cm, median (IQR)	165 (161-170)	163 (159-167)**
Maternal body mass index in kg/m <sup>2</sup> , median (IQR)	28.7 (25.9-32.3)	36.7 (27.4-34.4)**
Cigarette smoker, n (%)	2983 (8.6)	223 (8.6)
Racial origin		
White, n (%)	26216 (75.3)	1703 (65.8)
Black, n (%)	5213 (15.0)	597 (23.1)**
South Asian, n (%)	1612 (4.6)	160 (6.2)**
East Asian, n (%)	725 (2.1)	47 (1.8)
Mixed, n (%)	1068 (3.1)	83 (3.2)
Conception		
Natural, n (%)	33908 (97.3)	2469 (95.3)
Use of ovulation induction drugs, n (%)	177 (0.5)	17 (0.7)
<i>In vitro</i> fertilization, n (%)	749 (2.2)	104 (4.0)**
Obstetric history		
Nulliparous, n (%)	15464 (44.4)	1800 (69.5)**
Parous, previous SGA or PE, n (%)	3040 (8.7)	184 (7.1)**
Parous, no previous SGA or PE, n (%)	16330 (46.9)	606 (23.4)
Medical disorders		
Chronic hypertension, n (%)	328 (0.9)	55 (2.1)**
Diabetes mellitus, n (%)	189 (0.5)	30 (1.2)**
Pregnancy complications		
Gestational diabetes, n (%)	1208 (3.5)	128 (4.9)**
Obstetric cholestasis, n (%)	359 (1.0)	36 (1.4)
Doppler indices		
Umbilical artery PI in MoM, median (IQR)	1.01 (0.91-1.11)	1.02 (0.91-1.13)*
Umbilical artery PI >90 <sup>th</sup> percentile, n (%)	2975 (8.5)	257 (9.9)*
Middle cerebral artery PI in MoM, median (IQR)	1.00 (0.90-1.10)	0.98 (0.89-1.08)**
Middle cerebral artery PI <10 <sup>th</sup> percentile, n (%)	2822 (8.1)	270 (10.4)**
Cerebroplacental ratio in MoM, median (IQR)	0.99 (0.87-1.13)	0.97 (0.85-1.08)**
Cerebroplacental ratio <10 <sup>th</sup> percentile, n (%)	3307 (9.5)	340 (13.1)**
Estimated weight <10 <sup>th</sup> percentile, n (%)	3258 (9.4)	294 (11.4)**
GA at delivery in weeks, median (IQR)	40.1 (39.1-40.9)	40.4 (39.3-41.3)**
Birth weight in g, median (IQR)	3420 (3105-3730)	3350 (3000-3700)**
Birth weight <10 <sup>th</sup> percentile, n (%)	4040 (11.6)	503 (19.4)**

IQR = interquartile range; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median

Significance value \* p<0.05; \*\* p<0.01

**Table S5.** Univariable and multivariable logistic regression analysis in prediction of cesarean section from fetal distress from maternal and pregnancy characteristics.

Maternal and pregnancy characteristics	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Maternal age – 30 (years)	1.01 (1.00-1.02)	0.007	1.04 (1.03-1.04)	<0.0001
Maternal BMI – 30 (kg/m <sup>2</sup> )	1.06 (1.05-1.07)	<0.0001	1.07 (1.06-1.08)	<0.0001
Cigarette smoker	1.01 (0.87-1.16)	0.935	1.40 (1.20-1.63)	<0.0001
Racial origin				
White	1.00 (Reference)			
Black	1.76 (1.60-1.94)	<0.0001	1.90 (1.71-2.10)	<0.0001
South Asian	1.53 (1.29-1.81)	<0.0001	1.57 (1.32-1.87)	<0.0001
East Asian	1.00 (0.74-1.35)	0.989		
Mixed	1.20 (0.95-1.50)	0.124		
Conception				
Natural	1.00 (Reference)			
Use of ovulation induction drugs	1.32 (0.80-2.17)	0.277		
<i>In vitro</i> fertilization	1.91 (1.55-2.35)	<0.0001	1.34 (1.08-1.67)	0.009
Obstetric history				
Parous, no previous PE or SGA	1.00 (Reference)			
Nulliparous	3.14 (2.85-3.45)	<0.0001	3.92 (3.54-4.33)	<0.0001
Parous, previous PE or SGA	1.63 (1.38-1.93)	<0.0001	1.51 (1.27-1.80)	<0.0001
Medical complications				
Chronic hypertension	2.82 (1.71-3.05)	<0.0001	1.52 (1.13-2.06)	0.007
Diabetes mellitus	2.15 (1.46-3.16)	<0.0001	1.70 (1.14-2.54)	0.010
Pregnancy complications				
Gestational diabetes	1.45 (1.20-1.74)	<0.0001		
Cholestasis	1.35 (0.96-1.91)	0.085		
Estimated fetal weight				
Z-score	1.02 (0.98-1.06)	0.394		
<10 <sup>th</sup> percentile	1.24 (1.09-1.41)	0.001	1.25 (1.09-1.43)	0.001
Doppler indices				
UA-PI >90 <sup>th</sup> percentile	1.18 (1.03-1.35)	0.016		
MCA-PI <10 <sup>th</sup> percentile	1.32 (1.16-1.51)	<0.0001		
CPR <10 <sup>th</sup> percentile	1.44 (1.28-1.62)	<0.0001	1.31 (1.16-1.48)	<0.0001

BMI = body mass index; OR = odds ratio; CI = confidence interval; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median.

**Table S6.** Maternal and pregnancy characteristics in pregnancies delivering small for gestational age neonates with birthweight <3<sup>rd</sup> percentile compared to those with birthweight ≥3<sup>rd</sup> percentile.

Maternal and pregnancy characteristics	Birthweight ≥3 <sup>rd</sup> (n=45,109)	Birthweight <3 <sup>rd</sup> (n=2,102)
Maternal age in years, median (IQR)	31.6 (27.3-35.4)	31.0 (26.1-35.1)**
Maternal weight in kg, median (IQR)	79.0 (71.0-90.0)	73.0 (65.0-83.0)**
Maternal height in cm, median (IQR)	165 (161-169)	162 (158-167)**
Maternal body mass index in kg/m <sup>2</sup> , median (IQR)	29.2 (26.2-33.0)	27.7 (24.8-31.5)**
Cigarette smoker, n (%)	3480 (7.7)	371 (17.6)**
Racial origin		
White, n (%)	33795 (74.9)	1291 (61.4)**
Black, n (%)	7036 (15.6)	453 (21.6)**
South Asian, n (%)	2029 (4.5)	225 (10.7)**
East Asian, n (%)	917 (2.0)	51 (2.4)
Mixed, n (%)	1332 (3.0)	82 (3.9)*
Conception		
Natural, n (%)	43579 (96.6)	2013 (95.8)
Use of ovulation induction drugs, n (%)	252 (0.6)	12 (0.6)
<i>In vitro</i> fertilization, n (%)	1278 (2.8)	77 (3.7)*
Obstetric history		
Nulliparous, n (%)	20272 (44.9)	1193 (56.8)**
Parous, previous SGA or PE, n (%)	3798 (8.4)	4245 (20.2)**
Parous, no previous SGA or PE, n (%)	21039 (46.6)	485 (23.1)
Medical disorders		
Chronic hypertension, n (%)	560 (1.2)	36 (1.7)*
Diabetes mellitus, n (%)	372 (0.8)	9 (0.4)*
Pregnancy complications		
Gestational diabetes, n (%)	1936 (4.3)	97 (4.6)
Obstetric cholestasis, n (%)	483 (1.1)	15 (0.7)
Doppler indices		
Umbilical artery PI in MoM, median (IQR)	1.01 (0.91-1.11)	1.10 (0.99-1.20)**
Umbilical artery PI >90 <sup>th</sup> percentile, n (%)	3627 (8.0)	473 (22.5)**
Middle cerebral artery PI in MoM, median (IQR)	1.00 (0.90-1.10)	0.95 (0.86-1.04)**
Middle cerebral artery PI <10 <sup>th</sup> percentile, n (%)	3649 (8.1)	360 (17.1)**
Cerebroplacental ratio in MoM, median (IQR)	1.00 (0.89-1.13)	0.88 (0.77-1.01)**
Cerebroplacental ratio <10 <sup>th</sup> percentile, n (%)	4095 (9.1)	545 (25.9)**
Estimated weight <10 <sup>th</sup> percentile, n (%)	2927 (6.5)	1355 (64.5)
GA at delivery in weeks, median (IQR)	40.0 (39.0-40.9)	39.0 (37.9-40.0)**
Birth weight in g, median (IQR)	3450 (3150-3760)	2490 (2300-2635)**
Birth weight <10 <sup>th</sup> percentile, n (%)	3407 (7.6)	2102 (100.0)**

IQR = interquartile range; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median

Significance value \* p<0.05; \*\* p<0.01



**Table S7.** Univariable and multivariable logistic regression analysis in prediction of pregnancies delivering small for gestational age neonates with birthweight <3<sup>rd</sup> percentile from maternal and pregnancy characteristics.

Maternal and pregnancy characteristics	Univariate analysis		Multivariate analysis	
	OR (95% CI)	P value	OR (95% CI)	P value
Maternal age – 30 (years)	0.98 (0.97-0.99)	<0.0001	1.02 (1.01-1.03)	<0.0001
Maternal BMI – 30 (kg/m <sup>2</sup> )	0.94 (0.93-0.95)	<0.0001	0.98 (0.97-0.99)	0.001
Cigarette smoker	2.56 (2.28-2.88)	<0.0001	2.38 (2.05-2.76)	<0.0001
Racial origin				
White	1.00 (Reference)			
Black	1.69 (1.51-1.88)	<0.0001	2.04 (1.79-2.32)	<0.0001
South Asian	2.90 (2.50-3.36)	<0.0001	2.27 (1.90-2.71)	<0.0001
East Asian	1.46 (1.09-1.94)	0.010		
Mixed	1.61 (1.28-2.03)	<0.0001	1.72 (1.32-2.24)	<0.0001
Conception				
Natural	1.00 (Reference)			
Use of ovulation induction drugs	1.03 (0.58-1.84)	0.918		
<i>In vitro</i> fertilization	1.30 (1.03-1.65)	0.026		
Obstetric history				
Parous, no previous PE or SGA	1.00 (Reference)			
Nulliparous	2.55 (2.29-2.84)	<0.0001	2.34 (2.08-2.64)	<0.0001
Parous, previous PE or SGA	4.84 (4.23-5.54)	<0.0001	2.54 (2.18-2.97)	<0.0001
Medical complications				
Chronic hypertension	1.39 (0.99-1.95)	0.060		
Diabetes mellitus	0.52 (0.27-1.00)	0.051	0.44 (0.21-0.94)	0.033
Pregnancy complications				
Gestational diabetes	1.08 (0.88-1.33)	0.476		
Cholestasis	0.66 (0.39-1.11)	0.120		
Estimated fetal weight				
Z-score	0.15 (0.14-0.16)	<0.0001		
<10 <sup>th</sup> percentile	26.14 (23.72-28.8)	<0.0001	20.02 (18.07-22.19)	<0.0001
Doppler indices				
UA-PI >90 <sup>th</sup> percentile	3.32 (2.98-3.69)	<0.0001	1.67 (1.44-1.94)	<0.0001
MCA-PI <10 <sup>th</sup> percentile	2.35 (2.09-2.64)	<0.0001	1.58 (1.35-1.85)	<0.0001
CPR <10 <sup>th</sup> percentile	3.51 (3.16-3.89)	<0.0001	1.67 (1.43-1.95)	<0.0001

BMI = body mass index; OR = odds ratio; CI = confidence interval; SGA = small for gestational age with birthweight <10<sup>th</sup> percentile; PE = preeclampsia; PI = pulsatility index; MoM = multiple of the median.




Figure 1




Figure 2